

OPIC
OFFICE DE LA PROPRIÉTÉ
INTELLLECTUELLE DU CANADA



CIPO
CANADIAN INTELLECTUAL
PROPERTY OFFICE

(21) (A1) 2,132,766
(22) 1994/09/23
(43) 1995/04/08

(51) Int.Cl. ⁵
C07D-311/70; C07D-335/06; C07D-405/00; C07D-407/12;
C07D-413/12; C07D-413/14; C07D-417/12; C07D-417/14;
C07F-009/655; A61K-031/35; A61K-031/395; A61K-031/665

(19) (CA) **APPLICATION FOR CANADIAN PATENT** (12)

(54) 4-Arylamino-Benzopyran and Related Compounds

(72) Rovnyak, George C. - U.S.A. ;
Atwal, Karnail - U.S.A. ;
Santafianos, Dinos P. - U.S.A. ;
Ding, Charles Z. - U.S.A. ;

(71) Same as inventor

(30) (US) 134,034 1993/10/07

(57) 10 Claims

Notice: This application is as filed and may therefore contain an
incomplete specification.



Canada

2132766


HA641a

Abstract:

4-ARYLAMINO-BENZOPYRAN AND RELATED COMPOUNDS

Compounds having the formula

I



and pharmaceutically acceptable salts thereof wherein X is alkyl, Y is a single bond, $-\text{CH}_2-$, $-\text{C}(\text{O})-$, $-\text{O}-$, $-\text{S}-$ or $-\text{N}(\text{R}^8)-$ where R^8 is hydrogen, alkyl, haloalkyl, aryl, arylalkyl, cycloalkyl or (cycloalkyl)alkyl, and R^6 to R^7 are as defined herein. These compounds have potassium channel activating activity and are useful, therefore for example, as cardiovascular agents.


4-ARYLAMINO-BENZOPYRAN AND RELATED COMPOUNDS

This application is a continuation-in-part of U.S. application serial No. 08/134,034, filed October 7, 1993; incorporated by reference herein.

5

The present invention is directed to compounds of the formula

I




10 and pharmaceutically acceptable salts thereof. As used in formula I, and throughout the specification, the symbols have the following meanings:

a, b and d are all carbon atoms or one of a, b and d is a nitrogen atom or -N(O)- and the others are carbon atoms;



Y is a single bond, -CH₂-, -C(O)-, -O-, -S- or -N(R⁸)-;

15 R¹ is aryl or heterocyclo;

R² is -COOR⁸, -CO-amino, -CO-substituted amino, amino, substituted amino, -NR⁸CO-amino, -NR⁸CO-substituted amino, -NR⁸COR⁹, -NR⁸SO₂R⁹, -NR⁸(C=NCN)-amino, -NR⁸(C=NCN)-substituted amino,




$-\text{SR}^8$, $-\text{SOR}^8$, $-\text{SO}_2\text{R}^8$, $-\text{OR}^8$, cyano, heterocyclo, pyridine-N-oxide.



R^3 is hydrogen, hydroxy or $-\text{OC}(\text{O})\text{R}^8$;

5 R^4 and R^5 are each independently hydrogen, alkyl or arylalkyl, or R^4 and R^5 taken together with the carbon atom to which they are attached form a 5- to 7-membered carbocyclic ring;

R^6 is hydrogen, alkyl, haloalkyl, alkenyl, alkynyl, cycloalkyl, arylalkyl, (cycloalkyl)alkyl, $-\text{CN}$, $-\text{NO}_2$, $-\text{COR}^8$, $-\text{COOR}^8$,

10 $-\text{CONHR}^8$, $-\text{CONR}^8\text{R}^9$, $-\text{CF}_3$, $-\text{S-alkyl}$, $-\text{SOalkyl}$, $-\text{SO}_2\text{alkyl}$, $-\text{P}(\text{O})(\text{O-alkyl})_2$,  R^8 , halogen, amino, substituted amino, $-\text{O-alkyl}$, $-\text{OCF}_3$,

$-\text{OCH}_2\text{CF}_3$, $-\text{OCOalkyl}$, $-\text{OCONR}^8\text{alkyl}$, $-\text{NR}^8\text{COalkyl}$, $-\text{NR}^8\text{COOalkyl}$ or $-\text{NR}^8\text{CONR}^9$, tetrazolyl, imidazole, oxazole or triazole;

R^7 is hydrogen, alkyl, hydroxy, $-\text{O-alkyl}$, amino,

15 substituted amino, $-\text{NHCOR}^8$, $-\text{CN}$ or $-\text{NO}_2$;

R^8 and R^9 are independently hydrogen, alkyl, haloalkyl, aryl, arylalkyl, cycloalkyl or (cycloalkyl)alkyl;

X is alkyl; or X-R^2 together can be hydrogen, aryl or heterocyclo when R^1 is heterocyclo; and

20 n is an integer of 1 to 3.

The compounds of this invention possess antischismic activity and are useful, for example as cardiovascular agents.

25 The present invention provides for compounds of formula I, pharmaceutical compositions employing such compounds and for methods of using such compounds. Listed below are definitions of various terms used to describe the compounds of the instant invention. These definitions apply to the terms as they are used throughout the specification (unless

they are otherwise limited in specific instances either individually or as part of a larger group).

The term "alkyl" refers to both straight and branched chain groups having 1 to 8 carbon atoms preferably 1 to 5 carbons, such as methyl, ethyl, propyl, butyl, pentyl, hexyl, heptyl, octyl, the various branched chain isomers thereof, such as isopropyl, t-butyl, isobutyl, isohexyl, 4,4-dimethylpentyl, 2,2,4-trimethylpentyl and the like as well as such groups having a halo substituent (such as CCl_3 or CF_3), an alkoxy substituent, an aryl substituent, an alkylaryl substituent, a haloaryl substituent, a cycloalkyl substituent, a (cycloalkyl)alkyl substituent, a hydroxy substituent, an alkylamino substituent, an alkyl-substituted amino substituent, an alkanoylamino substituent, an arylcarbonylamino substituent, a nitro substituent, a cyano substituent, a thiol substituent or an alkylthio substituent.

The term "alkoxy" refers to any of the above alkyl groups linked to an oxygen atom.

The term "alkylthio" refers to any of the above alkyl groups linked to a sulfur atom.

The term "alkenyl" refers to any of the above alkyl groups further containing at least one carbon to carbon double bond.


The term "alkynyl" refers to any of the above alkyl groups further containing at least one carbon to carbon triple bond.

The term "cycloalkyl" refers to saturated cyclic hydrocarbon groups containing 3 to 7 ring carbons with cyclopropyl, cyclopentyl and cyclohexyl being preferred.

The term "halogen" or "halo" refers to chlorine, bromine, iodine and fluorine.

The term "aryl" refers to phenyl, 1-naphthyl, 2-naphthyl, phenyl, 1-naphthyl, 2-naphthyl, mono-substituted with $(\text{C}_1\text{-C}_4)$ -alkyl.

$(\text{C}_1\text{-C}_4)$ -alkylthio, $(\text{C}_1\text{-C}_4)$ -alkoxy, halo, nitro, cyano, hydroxy, amino, (alkyl)amino, alkyl-substituted amino, $-\text{NH}-(\text{C}_1\text{-C}_4)\text{-alkyl}$, $-\text{N}((\text{C}_1\text{-C}_4)\text{-alkyl})_2$, CF_3 , OCHF_2 , $-\text{OCH}_2-$, $-\text{SCH}_2-$



(where Z^1 is hydrogen, $(\text{C}_1\text{-C}_4)$ -alkyl, $(\text{C}_1\text{-C}_4)$ -alkylthio,

(C₁-C₄)-alkoxy, halo, hydroxy or -CF₃), -O-CH₂-cycloalkyl, or -S-CH₂-cycloalkyl; and phenyl, 1-naphthyl or 2-naphthyl, di-substituted with methyl, methoxy, methylthio, halo, -CF₃, nitro, amino, -OCHF₂, carboxylic acid or carboxylic ester. The term "aryl" also includes those groups listed above fused to a five- or six-membered ring which optionally contains an O, S or N atom (the nitrogen atom being substituted by an R⁷ group). Preferred aryl groups include unsubstituted phenyl and monosubstituted phenyl wherein the substituents are (C₁-C₄)-alkyl, methoxy, halo, nitro, cyano or -CF₃.

The term "heterocyclo" or "hetero" refers to fully saturated or unsaturated rings of 5 or 6 atoms containing one or two oxygen or sulphur atoms and/or one to four nitrogen atoms provided that the total number of hetero atoms in the ring is four or less. The hetero ring is attached by way of an available atom. Preferred monocyclic hetero groups include 2- and 3-thienyl, 2- and 3-furyl, 2-, 3- and 4-pyridyl and imidazolyl. The term "hetero" also includes bicyclic rings wherein the five- or six-membered ring containing oxygen or sulphur and/or nitrogen atoms as defined above is fused to a benzene ring and the bicyclic ring is attached by way of an available carbon atom. Preferred bicyclic hetero groups include 4-, 5-, 6- or 7-indolyl, 4-, 5-, 6- or 7-isindolyl, 5-, 6-, 7- or 8-quinolyl, 5-, 6-, 7- or 8-isouinolyl, 4-, 5-, 6- or 7-benzothiazolyl, 4-, 5-, 6- or 7-benzoxazolyl, 4-, 5-, 6- or 7-benzimidazolyl, 4-, 5-, 6- or 7-benzoxadiazolyl and 4-, 5-, 6- or 7-benzofurazanyl.

The term "heterocyclo" or "hetero" also includes such monocyclic and bicyclic rings wherein an available carbon atom is substituted with a (C₁-C₄)-alkyl, aryl, (C₁-C₄)-alkylthio, (C₁-C₄)-alkoxy, halo, nitro, keto, cyano, hydroxy, azo, thiazo, amino, -NH-(C₁-C₄)-alkyl, -N((C₁-C₄)-alkyl)₂, -CF₃, (aminoester)alkyl, carboxylic acid, carboxylic ester, -OCHF₂ or (C₁-C₄)-alkoxy further substituted with a carboxylic acid or such monocyclic and bicyclic rings wherein two or three available carbons have substituents selected from methyl, methoxy, methylthio, halo, -CF₃, nitro, hydroxy, amino and -OCHF₂.

The term "substituted amino" refers to a group of the formula -NZ²Z¹ wherein Z² is hydrogen, alkyl, cycloalkyl, aryl, arylalkyl,

(cycloalkyl)alkyl, morpholinylalkyl, heterocyclo or heterocycloalkyl and Z^3 is hydrogen, alkyl, cycloalkyl, aryl, arylalkyl, haloalkyl, hydroxyalkyl, alkoxyalkyl, thioalkyl, (cycloalkyl)alkyl or hydroxyalkyl further substituted with a carboxylic ester or carboxylic acid, with the proviso that when Z^2 is hydrogen, then Z^3 is other than hydrogen; or Z^2 and Z^3 taken together with the nitrogen atom to which they are attached are 1-pyrrolidinyl, 1-piperidinyl, 1-azepinyl, 4-morpholinyl, 4-thiomorpholinyl, 1-piperazinyl, 4-alkyl-1-piperazinyl, 4-arylalkyl-1-piperazinyl, 4-diarylalkyl-1-piperazinyl, 1-pyrrolidinyl, 1-piperidinyl, or 1-azepinyl, optionally substituted with alkyl, alkoxy, alkylthio, halo, trifluoromethyl or hydroxy.

The compounds of formula I can be present as salts, in particular pharmaceutically acceptable salts. If the compounds of formula I have, for example, at least one basic center, they can form acid addition salts. These are formed, for example, with strong inorganic acids, such as mineral acids, for example sulfuric acid, phosphoric acid or a hydrohalic acid, with strong organic carboxylic acids, such as alkanecarboxylic acids of 1 to 4 carbon atoms which are unsubstituted or substituted, for example, by halogen, for example acetic acid, such as saturated or unsaturated dicarboxylic acids, for example oxalic, malonic, succinic, maleic, fumaric, phthalic or terephthalic acid, such as hydroxycarboxylic acids, for example ascorbic, glycolic, lactic, malic, tartaric or citric acid, such as amino acids, (for example aspartic or glutamic acid or lysine or arginine), or benzoic acid, or with organic sulfonic acids, such as (C₁-C₄)-alkyl- or aryl-sulfonic acids which are unsubstituted or substituted, for example by halogen, for example methane- or p-toluene-sulfonic acid. Corresponding acid addition salts can also be formed having, if desired, an additionally present basic center. The compounds of formula I having at least one acid group (for example COOH) can also form salts with bases. Suitable salts with bases are, for example, metal salts, such as alkali metal or alkaline earth metal salts, for example sodium, potassium or magnesium salts, or salts with ammonia or an organic amine, such as morpholine, thiomorpholine, piperidine, pyrrolidine, a mono-, di- or tri lower alkylamine, for example ethyl-, tert-butyl-, diethyl-, diisopropyl-, triethyl-, tributyl- or dimethyl-

propylamine, or a mono-, di- or trihydroxy lower alkylamine, for example mono-, di- or triethanolamine. Corresponding internal salts may furthermore be formed. Salts which are unsuitable for pharmaceutical uses but which can be employed, for example, for the isolation or purification of free compounds I or their pharmaceutically acceptable salts, are also included.


Preferred salts of the compounds of formula I include monohydrochloride, hydrogensulfate, methanesulfonate, phosphate or nitrate.

All stereoisomers of the compounds of the instant invention are contemplated, either in admixture or in pure or substantially pure form. The compounds of the present invention can have asymmetric centers at any of the carbon atoms including any one of the R substituents. Consequently, compounds of formula I can exist in diastereomeric forms or in mixtures thereof. The above described processes can utilize racemates, enantiomers or diastereomers as starting materials. When diastereomeric products are prepared, they can be separated by conventional methods for example, chromatographic or fractional crystallization. Preferred compounds are those with the 3R or 4S stereochemistry.

It should be understood that the present invention includes prodrug forms of the compounds of formula I such as alkylesters of acids.

The compounds of the instant invention may, for example, be in the free or hydrate form, and may be obtained by methods exemplified by the following descriptions.

Compounds of formula I where R³ is trans-hydroxy and X is CH₂, can be prepared by first reacting an epoxide of formula II




with an amine of formula III



under heat or preferably in the presence of a Lewis acid such as magnesium perchlorate or trimethylaluminum to provide an intermediate of formula

IV




The intermediate of formula IV is then derivatized by reductive amination using an aldehyde of formula

V



in the presence of a reducing agent such as sodium cyanoborohydride or sodium triacetoxyborohydride. Alternatively, reductive amination can be effected with hydrogen gas in the presence of a catalyst such as palladium on carbon.

Compounds of formula I can also be prepared by reacting an epoxide of formula II with an amine of formula VI



in an organic solvent such as acetonitrile in the presence of a Lewis acid such as magnesium perchlorate or cobalt chloride.


Compounds of formula I wherein R² is CO-amino or CO-substituted amino, can be prepared by reacting compounds of formula I wherein R² is COOR⁶ with ammonia or an appropriate amine.

Compounds of formula I where R² is NR⁶CO-amino, NR⁶CO-substituted amino, NR⁶COR⁷, NR⁶SO₂R⁸, NR⁶(C=NCN)-amino or NR⁶(C=NCN)-substituted amino can be prepared from compounds of formula I where R² is amino or substituted amino by methods described in the literature such as those used for acylation, urea formation, sulfonylation and cyanoguanidine formation

Compounds of formula I where R^1 is heterocyclo (e.g., benzoxazole) and R^3 is trans-hydroxy can also be prepared by first reacting an epoxide of formula II with an amine of formula


VII

5



under heat or in the presence of a Lewis acid (magnesium perchlorate, trimethylaluminum, etc.) to provide an intermediate of formula

VIII



10

The intermediate of formula VIII is then reacted with a heterocycle containing a leaving group (e.g., 2-chlorobenzoxazole) in the presence of a base such as sodium hydride in an organic solvent such as tetrahydrofuran or dimethylformamide to form compounds of formula I where R^1 is heterocyclo and R^3 is trans-hydroxy.

15

Other compounds of formula I wherein R^1 is heterocyclo (e.g., oxazole, pyrazole, isonazole etc.) can be prepared from intermediates of formula VIII by standard methods.

Compounds of formula I wherein R^1 is heterocyclo (e.g., thiazole) can also be prepared by alkylation of a compound of formula IV with an

20

alkylating agent of formula

IX




where L is a leaving group such as a halogen, mesylate or tosylate.


Compounds of formula I wherein R^3 is hydrogen can be prepared from compounds of formula

25

X




by reaction with an amine of formula VI in the presence of a base such as sodium hydride or potassium carbonate. Alternatively, compounds of formula I where R³ is hydrogen can be prepared by first reacting a compound of formula X with an amine of formula III in the presence of a base (e.g., sodium hydride) to provide a compound of formula XI



The compound of formula XI is then converted to compounds of formula I where R³ is hydrogen by methods described for the conversion of compounds of formula IV to compounds of formula I.

Compounds of formula XI where R³ is hydrogen can also be prepared from the ketone of formula XII




and the amine of formula III by standard techniques of reductive amination.

The ketone of formula XII can be obtained by standard methodology or by literature procedures, such as those disclosed by P. Sebok and T. Timar, *Heterocycles*, 1988, 27, 2595; P. Teixidor et al., *Heterocycles*, 1988, 27, 2459; A. Benetti and N. C. Gooner, *Tetrahedron Letters*, 1979, 3685; and G. Ariamaia and K. K. Subramanian, *Tetrahedron Letters*, 1988, 29, No.28, 3487-3488.

The bromide of formula X can be prepared from the olefin of formula

XIII



by (a) catalytic hydrogenation of the double bond followed by (b) radical bromination using standard methods. The olefin of formula XIII can be prepared by the methods described for the preparation of compounds of formula II.

5 Compounds of formula XI where R^3 is hydrogen can also be prepared from compounds of formula IV by (a) dehydration of the alcohol with sodium hydride in aprotic solvents such as tetrahydrofuran; and (b) catalytic hydrogenation or reductive amination by sodium cyanoborohydride or sodium triacetoxyborohydride.

10 If any of the R substituents, X or Y groups contain reactive groups such as hydroxy or amino that can interfere with the epoxide opening reaction or any other reactions, they should be protected with appropriate protecting groups.

15 Compounds of formula II wherein Y is a single bond can be prepared according to D. R. Buckle, et al., J. Med. Chem., 1991, 34, 919.


 Compounds of formula II wherein Y is CH_2 can be prepared by methods described in V. A. Ashwood, et al., J. Med. Chem., 1991, 34, 3261.

20 Compounds of formula II where Y is oxygen, can be prepared by methods described in the literature, such as by J.M. Evans, et al., J. Med. Chem., 1983, 26, 1582; J.M. Evans, et al., J. Med. Chem., 1986, 29, 2194; R.W. Lang et al., Helvetica Chimica Acta, 1988, 71, 596; European patent 0205292 A2 and PCT patent 87/07607.

25 Compounds of formula II where Y is NiR can be prepared according to PCT patent 85/050083.


 To prepare enantiomers of epoxide II, the olefin of formula XIII is epoxidized with an oxidizing agent such as commercial bleach using a metal catalyst such as chiral manganese catalyst of the formula

XIV




as described by N.H. Lee, et al. (*Tetrahedron Letters*, 1991, 32,

5055-5058), to provide predominantly the chiral epoxide of formula
IIA



IIB




depending on the chirality of the 1,2-diaminocyclohexane used in the
preparation of a compound of formula XIV as described by Lee et al

The epoxides of formulae IIA and IIB can then be utilized to
prepare the chiral compounds of formula I

Compounds of formula I where R¹ is OC(O)R⁶ can be prepared
from compounds of formula I where R¹ is hydroxy by treatment with an
acid chloride of formula

XV



in the presence of a base catalyst such as pyridine or triethylamine.

All other compounds of formula I may be prepared by
modification of the procedures discussed herein as known by those skilled

in the art. The intermediates used to prepare compounds of formula I are described herein or may be derived from known compounds by those skilled in the art or are commercially available.

The compounds of the present invention can have asymmetric centers at carbons 2-4 of the bicyclic ring. Also, any one of the R's can have an asymmetric carbon. Consequently, compounds of formula I can exist in diastereomeric forms or in mixtures thereof. The above described process can utilize racemates, enantiomers or diastereomers as starting materials. When diastereomeric products are prepared, they can be separated by conventional chromatographic or fractional crystallization methods.

The antiischemic and antihypertensive effects of benzopyran based and related potassium channel openers are usually stereoselective, with the 3S,4R- enantiomer being the more active isomer. However, it has been unexpectedly found that compounds of formula I are "selective antiischemic agents" with the 3R,4S-enantiomer being the more potent isomer. The term "selective antiischemic agent" means that these compounds possess little or no vasodilator activity (i.e., these compounds have IC₅₀ (rat aorta) values greater than that of the known potassium channel activator, cromakalim. Therefore, in the treatment of ischemic hearts, the compounds of the instant invention are less likely to cause coronary steal, profound hypotension and coronary under-perfusion.

The preferred compounds of the present invention are those compounds of formula I where:

- a, b and d are carbon atoms;
- X is alkyl;
- Y is a single bond or -(O)-;
- R¹ is aryl or heterocyclo;
- R² is -COOR⁴, -CO-amino, -CO-substituted amino,
- NHCOCH₃, -NHSO₂Me, -NHCONH₂, -NHC≡NCN(NH₂), imidazole,
- 30 furan, pyridine, oxazole, hydroxy, -NHCO-substituted amino or -SO₂Me;
- R³ is hydroxy,
- R⁴ and R⁵ are methyl,
- R⁶ is cyano, -NO₂, -CF₃, halo, alkyl or tetrazol; and

R⁷ is hydrogen.

Compounds of formula I may be used as antiischemic agents, i.e., for the treatment of ischemic conditions such as myocardial ischemia, cerebral ischemia, lower limb ischemia and the like.

5 Thus a composition containing one (or a combination) of the compounds of this invention, may be administered to a species of mammal (e.g., humans) suffering from an ischemic or hypertensive condition.

A single dose, or two to four divided daily doses, provided on a basis of about 0.001 to about 100 mg per kilogram of body weight per day, preferably about 0.1 to about 25 mg per kilogram of body weight per day
10 is appropriate. The substance is preferably administered orally, but parenteral routes such as the subcutaneous, intramuscular, intravenous or intraperitoneal routes or any other suitable delivery system, such as intranasal or transdermal routes can also be employed.

15 As a result of the potassium channel activating activity of the compounds of this invention, these compounds are also useful in the treatment of cardiovascular disorders and any disorders associated with smooth muscle contraction. For example, compounds of the present invention are useful as therapy for congestive heart failure, therapy for
20 peripheral vascular disorders (e.g. Raynaud's Disease), therapy for pulmonary hypertension, as anti-anginal agents, as antifibrillatory agents, and in limiting myocardial infarction.

Compounds of the present invention are additionally expected to be useful in the treatment of central nervous system disorders (e.g.,
25 Parkinsonism, as anti-tremor agents, epilepsy), in therapy for renal failure, in therapy for urinary incontinence, as anti-diarrheal agents, in therapy for pre-eclampsia, dysmenorrhea and premature labor, for the treatment of male impotence, as well as for the promotion of hair growth (e.g., in the treatment of male pattern baldness), and as anti-asthmatic agents.

30 The compounds of this invention can also be formulated in combination with a diuretic such as chlorothiazide, hydrochlorothiazide.


musolimine, bumetanide, triamterene, amiloride and spironolactone and salts of such compounds, angiotensin converting enzyme inhibitors such as captopril, zofenopril, fosinopril, enalapril, ceranopril, cilazapril, delapril, pentopril, quinapril, ramipril, lisinopril, and salts of such compounds, thrombolytic agents such as tissue plasminogen activator (tPA), recombinant tPA, streptokinase, urokinase, procurokinase, and anisoylated plasminogen streptokinase activator complex (APSAC, Eminase, Beecham Laboratories), or calcium channel blocking agents such as nifedipine or diltiazem. Such combination products if formulated as a fixed dose employ the compounds of this invention within the dose range described above and the other pharmaceutically active agent within its approved dose range.

The compounds of formula I, and combinations thereof, can be formulated, as described above, in compositions such as tablets, capsules or elixirs for oral administration, in sterile solutions or suspensions for parenteral administration, and may also be administered via transdermal patch or nasal inhalation solutions. About 10 to about 500 milligrams of a compound of formula I is compounded with physiologically acceptable vehicle, carrier, excipient, binder, preservative, stabilizer, flavor, etc., in a unit dosage form as called for by accepted pharmaceutical practice. The amount of active substance in these compositions or preparations is such that a suitable dosage in the range indicated is obtained.

The following examples and preparations describe the manner and process of making and using the invention and are illustrative rather than limiting. It should be understood that there may be other embodiments which fall within the spirit and scope of the invention as defined by the claims appended hereto.

Example 1

trans-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid, ethyl ester



A solution of 3,4-dihydro-2,2-dimethyl-3,4-epoxy-2H-benzopyran-6-carbonitrile (400 mg, 2.0 mmol, prepared according to Evans et al., *J. Med. Chem.*, 1983, 26, 1582 and *J. Med. Chem.*, 1986, 29, 2194) and *D*-phenylglycine ethyl ester (700 mg, 4.0 mmol) in acetonitrile (10 mL) under argon at room temperature was treated with magnesium perchlorate (450 mg, 2.0 mmol). The mixture was stirred at 45°C for two days; diluted with ethyl acetate and washed with 5% sodium bicarbonate, water and brine. The dried (anhydrous magnesium sulfate) organic solution was concentrated to give an oil. Flash chromatography on silica gel eluting with ethyl acetate/hexanes (1:10) gave a foam (450 mg). Trituration with hexanes gave the title product (400 mg, 53%) as a colorless solid, mp 140-144°C. Analysis calculated for $C_{27}H_{24}N_2O_4$: C, 69.46; H, 6.36; N, 7.36. Found: C, 69.22; H, 6.37; N, 7.28.

Example 2

(3S-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid, ethyl ester



A. (1aS-cis)-1a,7b-Dihydro-2,2-dimethyl-2H-oxireno-
[c][1]benzopyran-6-carbonitrile


The title compound was prepared by the procedure described by Lee et
al., *Tetrahedron Letters*, 1991, 32, 5055.

B. (3S-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-
benzopyran-4-yl)phenylamino]acetic acid, ethyl ester

The title compound was prepared from the title A compound and N-
phenylglycine ethyl ester by the same procedure as described in Example
1. The product was purified by flash chromatography on silica gel eluting
with ethyl acetate/hexanes (1:10) to give a foam which was triturated with
hexanes to give the title compound as a colorless solid, mp 182-183°C.
Analysis calculated for $C_{22}H_{24}N_2O_4 \cdot 0.24 H_2O$: C, 68.69; H, 6.41; N, 7.28.
Found: C, 68.57; H, 6.28; N, 7.40. $[\alpha]_D = -100.3$ (c = 1.08, $CDCl_3$).

Example 3

(3R-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-
benzopyran-4-yl)phenylamino]acetic acid, ethyl ester



A. (1aR-cis)-1a,7b-Dihydro-2,2-dimethyl-2H-oxireno-
[c][1]benzopyran-6-carbonitrile

The title compound was prepared by the procedure described by Lee et
al., *Tetrahedron Letters*, 1991, 32, 5055.


B. (3R-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid, ethyl ester

The title compound was prepared from the title A compound and N-phenylglycine ethyl ester by the same procedure as described in Example 1.

1. The product was obtained as a colorless solid, mp 182-183 °C. Analysis calculated for $C_{22}H_{24}N_2O_4 \cdot 0.37 H_2O$: C, 68.26; H, 6.44; N, 7.24. Found: C, 67.91; H, 6.04; N, 7.59. $[\alpha]_D^{25} = +97.2$ ($c = 0.82$, $CDCl_3$).

Example 4


trans-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid



- 15 A solution of trans-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid, ethyl ester (310 mg, 0.81 mmol, the title compound of Example 1) in tetrahydrofuran (7 mL) and water (5 mL) at 0-5 °C was treated with 1M lithium hydroxide (1 mL) and stirred for three hours as the temperature rose to ambient. The mixture was diluted with ethyl acetate and extracted with water (2x). The combined aqueous fractions were acidified with 10% citric acid to pH 3 and extracted with ethyl acetate. The organic fraction was washed with water and brine, dried (anhydrous magnesium sulfate) and concentrated *in vacuo* to give the product as a foam. Trituration with hexane containing 1-2% ether afforded the title product (230 mg) as a colorless solid, mp 163-165 °C. Analysis calculated for $C_{20}H_{20}N_2O_4 \cdot 0.11 H_2O$: C, 67.79; H, 5.75; N, 7.91. Found: C, 67.70; H, 5.66; N, 8.00.

Example 5

(3R-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-fluorophenyl)amino]acetic acid, ethyl ester



A. N-(4-Fluorophenyl)glycine ethyl ester


A solution of ethyl glyoxylate (2.47 g, 0.024 mol) in 1,2-dichloroethane (30 mL) under argon at room temperature was treated successively with 4-fluoroaniline (1.80 g, 0.016 mol), sodium triacetoxyborohydride (5.12 g, 0.024 mol) and acetic acid (1 mL). After stirring for two hours, the mixture was concentrated, dissolved in ethyl acetate and washed 5% sodium bicarbonate, water and brine. The dried (anhydrous magnesium sulfate) organic solution was concentrated and crystallized from ether/hexanes to give the title product (2.07 g, 65%) as a colorless solid, mp 72-73°C. Analysis calculated for $C_{10}H_{12}FNO_2 \cdot 0.07 H_2O$: C, 60.54; H, 6.16; N, 7.06; F, 9.58. Found: C, 60.75; H, 6.15; N, 6.96; F, 9.13.

B. (3R-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-fluorophenyl)amino]acetic acid, ethyl ester

The title compound was prepared from the title A compound and (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) by the same procedure as described in Example 1. The product was purified by flash chromatography on silica gel, eluting with ethyl acetate/hexanes (1:12) to give a foam which was crystallized from ethyl acetate/hexanes to provide the title compound as a colorless solid (297 mg, 37%); mp 195-197°C. Analysis calculated for $C_{22}H_{23}FN_2O_4$: C, 66.32; H, 5.82; N, 7.03; F, 4.77. Found: C, 66.25; H, 5.78; N, 7.03; F, 4.87. $[\alpha]_D^{25} = +60.0^\circ$ (c = 0.58, $CHCl_3$).

Example 6

(3R-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-chlorophenyl)amino]acetic acid, ethyl ester



A. N-(4-Chlorophenyl)glycine ethyl ester


The title compound was prepared by the same procedure as described in Example 5, part A. The product was crystallized from ether/hexanes to give a colorless solid (1.71 g, 52%), mp 93-95°C. Analysis calculated for $C_{10}H_{12}ClNO_2$: C, 56.21; H, 5.66; N, 6.56; Cl, 16.59. Found: C, 56.10; H, 5.65; N, 6.44; Cl, 16.78.

B. (3R-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-chlorophenyl)amino]acetic acid, ethyl ester

To a mixture of (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (200 mg, 1.0 mmol, the title A compound of Example 3), N-(4-chlorophenyl)glycine ethyl ester (250 mg, 1.17 mmol, the title A compound) and magnesium perchlorate (225 mg, 1.0 mmol) under argon at room temperature was added acetonitrile (0.4 mL). The mixture was stirred at room temperature for three days. The ethyl acetate diluted solution was adsorbed onto celite and flash chromatographed on silica gel, eluting with ethyl acetate/hexanes (1:12) to give the product as a foam (210 mg). Trituration with hexanes gave the title compound (195 mg, 47%), mp 171.5-171.5°C. Analysis calculated for $C_{22}H_{23}ClN_2O_4$: C, 63.69; H, 5.59; N, 6.75; Cl, 8.55. Found: C, 63.52; H, 5.43; N, 6.43; Cl, 8.26. $[\alpha]_D^{25} = +105.2^\circ$ (c = 0.40, $CDCl_3$).

Example 7

(3R-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetamide




5

A mixture of (3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid, ethyl ester (310 mg, 0.81 mmol, the title compound of Example 3) in methanol (1 mL) was
 10 treated with 9.5 M methanolic ammonia (2 mL). After stirring at room temperature for 48 hours, volatiles were removed *in vacuo* to give a solid which was triturated with hexanes to afford title compound (286 mg, 100%) as a colorless solid, mp 236-238 °C. Analysis calculated for $C_{20}H_{21}N_3O_3 \cdot 0.22 H_2O$: C, 67.60; H, 6.08; N, 11.82. Found: C, 67.65; H, 6.05; N, 11.77. $[\alpha]_D^{25} = +90.2^\circ$ ($c = 1.14$, 10:1 $CDCl_3/CH_3CN$).
 15

Example 8

(3S-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[(4-phenyl-2-thiazolyl)amino]-2H-1-benzopyran-6-carbonitrile

20



The title compound was prepared from 2-amino-4-phenylthiazole and (1aS-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno[6,1]benzopyran-6-carbonitrile (the title A compound of Example 2), by the procedure described in Example 1. The residue was purified by flash
 25

2132766


HA641a

- 21 -

chromatography on silica gel (25% ethyl acetate in hexanes) to yield a light yellow foam which was crystallized from ether-hexanes to provide the title compound as a light yellow solid, mp 203-204°C. Analysis calculated for $C_{21}H_{19}N_3O_2S$: C, 66.82; H, 5.07; N, 11.13; S, 8.49. Found: C, 66.83; H, 5.14; N, 10.98; S, 8.54. $[\alpha]_D = -31.4^\circ$ (c = 0.5, MeOH).

Example 9


(3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[(4-phenyl-2-thiazolyl)amino]-2H-1-benzopyran-6-carbonitrile



The title compound was prepared from 2-amino-4-phenylthiazole and (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3), by the procedure described in Example 1 to give the title compound as a light yellow solid, mp 200-202°C. Analysis calculated for $C_{21}H_{19}N_3O$: S: C, 66.82; H, 5.07; N, 11.13; S, 8.49. Found: C, 66.61; H, 5.12; N, 10.94; S, 8.64. $[\alpha]_D = +31.7^\circ$ (c = 0.5, MeOH).

Example 10


(3R-trans)-[N-[3,4-Dihydro-3-hydroxy-2,2-dimethyl-6-(1H-tetrazol-5-yl)-2H-1-benzopyran-4-yl]phenylamino]acetic acid, ethyl ester



A mixture of (3R-trans)-[6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl]phenylamino]acetic acid, ethyl ester (420 mg, 1.1 mmol, the title compound of Example 3), sodium azide (190 mg, 3.0 mmol) and ammonium chloride (150 mg, 3.0 mmol) in dimethylformamide (1 mL) under argon was heated at 85°C for two days. The reaction mixture was then poured into water (50 mL), extracted with ethyl acetate (3x100 mL) and washed with water (3x100 mL). After drying over anhydrous magnesium sulfate, the solvent was evaporated and the residue was purified by flash chromatography (5% methanol in dichloromethane) to give the title product (300 mg, 64%). This material was combined with another batch of the same product and rechromatographed on silica gel (5% methanol in dichloromethane). The product was recrystallized from isopropyl ether-hexanes to give (3R-trans)-[N-[3,4-dihydro-3-hydroxy-2,2-dimethyl-6-(1H-tetrazol-5-yl)-2H-1-benzopyran-4-yl]phenylamino]acetic acid, ethyl ester, mp 130-133°C. Analysis calculated for $C_{22}H_{25}N_5O_4 \cdot 0.26H_2O$: C, 61.71; H, 6.01; N, 16.35. Found: C, 61.85; H, 6.13; N, 16.21. $[\alpha]_D^{25} = +92.4^\circ$ ($c = 0.392$, $CDCl_3$).

Example 11

(3R-trans)-2-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-ethylacetamide



5


The title compound was prepared from (3R-trans)-[6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl]phenylamino]acetic acid, ethyl ester (the title compound of Example 3) and ethyl amine by the same procedure as described for the title compound of Example 7. The product was triturated with hexanes to give the title compound as a colorless solid, mp 213-215°C. Analysis calculated for $C_{22}H_{25}N_3O_3$: C, 69.64; H, 6.64; N, 11.07. Found: C, 69.31; H, 6.33; N, 10.96. $[\alpha]_D^{25} = +76.6^\circ$ (c = 0.47, $CDCl_3$).

15

Example 12

(3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N[2-(1-pyrrolidinyl)-2-oxoethyl]phenylamino]-2H-1-benzopyran-6-carbonitrile

20




The title compound was prepared from (3R-trans)-[6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl]phenylamino]acetic acid,

25

ethyl ester (the title compound of Example 3) and pyrrolidine by the same procedure as described for the title compound of Example 7. The product was triturated with hexanes to give the title compound as a colorless solid, m.p. 222-224°C. Analysis calculated for $C_{24}H_{27}N_3O_3 \cdot 0.17H_2O$: C, 70.55; H, 6.75; N, 10.28. Found: C, 70.61; H, 6.76; N, 10.22. $[\alpha]_D^{25} = +45.6^\circ$ (c = 0.78, DMSO).

Example 13


(3R-4trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N[2-(4-morpholinyl)-2-oxoethyl]phenylamino]-2H-1-benzopyran-6-carbonitrile



The title compound was prepared from (3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid, ethyl ester (the title compound of Example 3) and morpholine by the same procedure as described for the title compound of Example 7. The product was triturated with hexanes to give the title compound as a colorless solid, mp 229-231°C. Analysis calculated for $C_{24}H_{27}N_3O_4 \cdot 0.07H_2O$: C, 68.17; H, 6.47; N, 9.94. Found: C, 68.29; H, 6.46; N, 9.82. $[\alpha]_D^{25} = +54.6^\circ$ (c = 0.71, DMSO).

Example 14

(3R-trans)-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-(2-furanylmethyl)acetamide



5


The title compound was prepared from (3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid ethyl ester (the title compound of Example 3) and 2-furanylmethyl amine by the same procedure as described for the title compound of Example 7. The product was obtained as a colorless solid, mp 95-100°C. Analysis: calculated for $C_{25}H_{25}N_3O_4 \cdot 0.25H_2O$: C, 68.89; H, 5.89; N, 9.64. Found: C, 68.73; H, 5.98; N, 9.42. $[\alpha]_D^{25} = +26.5^\circ$ ($c = 0.29$, MeOH).

10

15

Example 15

(3R-trans)-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-[2-(4-morpholinyl)ethyl]acetamide




20

The title compound was prepared from (3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid ethyl ester (the title compound of Example 3) and 4-morpholineethyl

amine by the same procedure as described for the title compound of Example 7. The product was obtained as a colorless solid, mp 201-204°C. Analysis calculated for $C_{26}H_{32}N_4O_4$: C, 67.22; H, 6.94; N, 12.06. Found: C, 67.09; H, 5.88; N, 11.88. $[\alpha]_D = +23.7^\circ$ (c = 0.43, MeOH).

Example 16

(3R-trans)-4-[(4-Fluorophenyl)(2-hydroxy-2-methylpropyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



A. N-(4-Fluorophenyl)-N-(2-hydroxy-2-methylpropyl)amine

A mixture of 4-fluoroaniline (1.12 g, 10 mmol) and isobutene oxide (0.70 g, 10 mmol) was heated in a sealed tube at 120°C overnight. The resultant oil was purified by flash chromatography to give the title compound as an oil (1.20 g, 55%).


B. (3R-trans)-4-[(4-Fluorophenyl)(2-hydroxy-2-methylpropyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

The title compound was prepared from title A compound and (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) by the procedure described in

Example 1. The product was obtained as an amorphous solid, mp 70°C. $[\alpha]_D = -62.8^\circ$ (c = 1, CHCl₃). Analysis calculated for $C_{22}H_{24}N_2O_4$ (f(0.3 H₂O=0.2 toluene): C, 68.84; H, 6.71; N, 6.86. Found: C, 68.84; H, 6.72; N, 6.59.

Example 17

[3R-{3a,4b(R*)}] 4-[(4-Fluorophenyl)(2-hydroxypropyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



A. N-(4-Fluorophenyl)-N-(2-hydroxypropyl)amine


The title compound was prepared from 4-fluoroaniline and R-propylene oxide by the same procedure as described for the title A compound of Example 15. The product was obtained as a colorless solid.

B. [3R-{3a,4b(R*)}] 4-[(4-Fluorophenyl)(2-hydroxypropyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

The title compound was prepared from title A compound and (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) by the procedure described for the title compound of Example 1. The product was obtained as an amorphous solid, mp 75°C. $[\alpha]_D^{25} = -64.2^\circ$ (c = 1, CHCl₃). Analysis calculated for C₂₁H₂₃N₂O₃F·0.73 H₂O: C, 65.59; H, 6.44; N, 7.29. Found: C, 65.40; H, 6.30; N, 7.48.

Example 18

(3R-*trans*)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-methyl-2-thiazolyl)amino]acetic acid, ethyl ester



A. (3R-*trans*)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[(4-methyl-2-thiazolyl)amino]-2H-1-benzopyran-6-carbonitrile

The title compound was prepared from (1aR-*cis*)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) and 2-amino-4-methyl-thiazole by the procedure described for the title compound of Example 1. The residue was purified by a flash column to give a colorless solid (730 mg, 53%).

B. (3R-*trans*)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-methyl-2-thiazolyl)amino]acetic acid, ethyl ester

A solution of title A compound (620 mg, 2.0 mmol) and ethyl bromoacetate (0.24 mL, 2.15 mmol) in dimethylformamide (4 mL) was treated with potassium carbonate (300 mg, 2.17 mmol). The resultant reaction mixture was stirred at room temperature overnight and poured into saturated sodium bicarbonate (10 mL). The aqueous solution was extracted with ethyl acetate (2 x 40 mL); combined organic extracts were dried over sodium sulfate, and concentrated *in vacuo*. The residue was purified by flash column chromatography to give an oil which solidified upon vacuum drying. The solid was washed with pentane to provide a colorless product (110 mg, 14%), mp 83°C. $[\alpha]_D^{25} = +33.2^\circ$ (c = 1, MeOH). Analysis calculated for $C_{20}H_{23}N_2SO_4 \cdot 1.31 H_2O$: C, 56.51; H, 6.07; N, 9.89. Found: C, 56.81; H, 5.96; N, 9.59.

Example 19

(3R-trans)-4-[N-(2-Benzoxazolyl)-N-(2,2-dimethoxyethylamino)]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



A. (3R-trans)-3,4-Dihydro-4-[(2,2-dimethoxyethylamino)-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

A mixture of (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (750 mg, 3.73 mmol, the title A compound of Example 3), dimethyl aminoacetal (1.2 mL) was heated in a sealed tube at 75°C for two days. The reaction mixture was purified by flash column chromatography (ethyl acetate and hexane, 1:1) to give a colorless oil (1.0 g, 90%).

B. (3R-trans)-4-[N-(2-Benzoxazolyl)-N-(2,2-dimethoxyethyl)-amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

To a solution of title A compound (360 mg, 1.18 mmol) in dimethylformamide (10 mL) at 0°C under argon was added sodium hydride (60% in oil, 100 mg, 2.3 mmol). The suspension was stirred at 0°C for 15 minutes and then treated with 2-chlorobenzoxazole (140 µL, 1.18 mmol) via a syringe. The reaction mixture was stirred at 0°C for 30 minutes and poured into saturated ammonium chloride. The aqueous solution was extracted with ethyl acetate; the combined organic extracts were treated with acetic acid (0.5 mL) and stirred at room temperature overnight. The resultant solution was washed with sodium bicarbonate, dried over sodium sulfate and concentrated *in vacuo*. The residue was purified by flash column chromatography (3:1 mixture of hexane and ethyl acetate) to give an oil, which formed a foam upon vacuum drying.


(300 mg, 60%), mp 66°C. $[\alpha]_D^{25} = +35.8^\circ$ ($c = 1$, MeOH). Analysis calculated for $C_{23}H_{25}N_3O_5 \cdot 0.33 H_2O \cdot 0.40$ toluene: C, 66.46; H, 6.24; N, 9.01. Found: C, 66.47; H, 6.21; N, 8.81.

5

Example 20

(3R-trans)-4-[N-[1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)ethyl]-phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

10



A. N-[1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)ethyl]-N-phenylamine

- 15 A solution of N-phenylethylenediamine (10.0 g, 73.4 mmol) and phthalic anhydride (11.42 g, 77.1 mmol) in toluene was heated at reflux while removing water azeotropically for 18 hours. The reaction mixture was cooled to room temperature and washed with 2.5% hydrochloric acid solution, saturated sodium bicarbonate solution and brine. The crude product solution was dried over magnesium sulfate and the solvent was recovered under vacuum to provide the title product (12.03 g, 61%) as a yellow solid, mp 100-102°C. MS: $(M+NH_4)^+$ @ 267.
- 20


B. (3R-trans)-4-[N-[1,3-Dihydro-1,3-dioxo-2H-isoindol-2-yl)ethyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

- 25 The title compound was prepared from title A compound (6.62 g, 24.85 mmol) and (1R-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (5.0 g, 24.85 mmol, the title A compound

- of Example 3) by the procedure described for the title compound of Example 1. The crude product was purified by flash chromatography on silica gel eluting with hexane/acetone (3:1) to afford a yellow solid (7.87 g, 67%), mp 150-160°C. $[\alpha]_D = -110.0^\circ$ ($c = 0.88$, CHCl_3).
- 5 Analysis calculated for $\text{C}_{28}\text{H}_{25}\text{N}_3\text{O}_4 \cdot 0.16 \text{ H}_2\text{O}$: C, 71.49; H, 5.43; N, 8.93. Found: C, 71.54; H, 5.49; N, 8.88.

Example 21


- 10 (3R-trans)-4-[N-(2-Aminoethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



- 15 To a solution of the title compound of Example 20 (4.5 g, 10.69 mmol) in ethanol (100 mL) at room temperature was added a mixture of methyl hydrazine (50 mL) and ethanol (50 mL). The reaction was stirred at room temperature for 1.5 hours and heated at reflux for one hour. The volatiles were recovered under vacuum and the residue was partitioned between
- 20 ethyl acetate and saturated sodium bicarbonate solution. The organic phase was washed with brine, dried over magnesium sulfate and evaporated *in vacuo* to obtain an off-white foam (3.57 g, 100%) which was crystallized from isopropyl ether/ethyl acetate to provide a colorless solid, mp 164-165°C. $[\alpha]_D = -20.9^\circ$ ($c = 1.22$, CHCl_3). Analysis
- 25 calculated for $\text{C}_{20}\text{H}_{23}\text{N}_3\text{O}_2 \cdot 1.0 \text{ H}_2\text{O} \cdot 0.42 \text{ isopropyl ether}$: C, 67.90; H, 7.81; N, 10.55. Found: C, 67.90; H, 7.38; N, 10.18.

Example 22

(3R-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]butyric acid, ethyl ester



5


The title compound was prepared from 4-(N-phenylamino)-butyric acid (prepared according to literature methods) and (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) by the procedure described for the title compound of Example 1. The product was triturated with hexanes to yield an off-white solid, mp 109-110°C. $[\alpha]_D = +41.8^\circ$ (c = 0.92, MeOH). Analysis calculated for $C_{24}H_{28}N_2O_4 \cdot 0.09 H_2O$: C, 70.28; H, 6.93; N, 6.83. Found: C, 70.31; H, 6.81; N, 6.80.

15

Example 23

(3R-trans)-3-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]propanoic acid, ethyl ester

20



The title compound was prepared from 3-(N-phenylamino)-propanoic acid (prepared according to literature methods) and (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) by the procedure described for the title compound of Example 1. The product was triturated with hexanes to yield


25

an off-white solid. mp 60-62°C. $[\alpha]_D^{25} = +30.4^\circ$ ($c = 0.8$, MeOH).
 Analysis calculated for $C_{23}H_{26}N_2O_4 \cdot 0.1 H_2O$: C, 69.70; H, 6.66; N, 7.07.
 Found: C, 69.75; H, 6.74; N, 7.02.

5

Example 24

(3R-*trans*)-3,4-Dihydro-3-hydroxy-4-[N-[(1*H*-imidazol-2-yl)-
 methyl]phenylamino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



10

A. N-[(1*H*-Imidazol-2-yl)methyl]-*N*-phenylamine


A stirred solution of 2-imidazolecarboxaldehyde (400 mg, 4.16 mmol) in
 dry methanol (10 mL) was treated with aniline (380 μ L, 4.16 mmol)
 15 followed by anhydrous magnesium sulfate (2 g). The solution was stirred
 for 18 hours at room temperature. The mixture was filtered and most of
 the solvent was evaporated. The solution was taken up in ethyl acetate
 (50 mL) and washed with 5% aqueous $KHSO_4$ (50 mL) followed by brine
 (50 mL). The organic solution was dried ($MgSO_4$), filtered and solvent
 20 was removed *in vacuo* to give a white solid (370 mg, 52%). This material
 (370 mg, 2.16 mmol) in MeOH (10 mL) was treated with 10% palladium
 over charcoal and hydrogenated at room temperature using a balloon. The
 mixture was filtered through a short pad of celite and the solvent volume
 was removed *in vacuo* to give a light brown crystalline solid (350 mg,
 25 93%). Analysis calculated for $C_{13}H_{10}N_2 \cdot 0.07 H_2O$: C, 68.82; H, 6.44,
 N, 24.08. Found: C, 68.77; H, 6.47; N, 24.13.

B. (3R-trans)-3,4-Dihydro-3-hydroxy-4-[N-[(1H-imidazol-2-yl)-methyl]phenylamino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

- A stirred solution of (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (792 mg, 3.93 mmol, the title A compound of Example 3) in acetonitrile (4 mL) was treated with the title A compound (620 mg, 3.62 mmol) followed by anhydrous COCl_2 (46.5 mg, 0.36 mmol). The solution was stirred for 18 hours at room temperature. Additional COCl_2 was added over an 18-hour period. The mixture was added to ethyl acetate (100 mL) and water (100 mL). The organic fraction was washed with brine (100 mL), dried over magnesium sulfate and the solvent was removed *in vacuo* to give a white solid which was purified by flash chromatography on silica gel (25% ethyl acetate in hexanes). The product was re-crystallized from chloroform-hexanes to give a colorless solid (321 mg, 24%), mp 259-260°C (shrinks at 140-150°C). $[\alpha]_D^{25} = +16.5^\circ$ (c = 0.31, MeOH). Analysis calculated for $\text{C}_{22}\text{H}_{22}\text{N}_4\text{O}_2 \cdot 0.94 \text{ H}_2\text{O}$: C, 67.51; H, 6.15; N, 14.32. Found: C, 67.75; H, 5.82; N, 14.08.

Example 25


- 20 (3R-trans)-4-[(2-(Acetylamino)ethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



- 25 The title compound was prepared from the title compound of Example 20 by a standard procedure using acetyl chloride in pyridine and dichloromethane. The product was obtained as a colorless solid, mp 217-218°C. $[\alpha]_D^{25} = +54.6^\circ$ (c = 1.15, DMF). Analysis calculated for $\text{C}_{22}\text{H}_{25}\text{N}_3\text{O}_4 \cdot 21 \text{ H}_2\text{O}$: C, 68.95; H, 6.68; N, 10.97. Found: C, 69.19; H, 6.74; N, 10.73.

Example 26


(3R-trans)-[2-(N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino)ethyl]urea



The title compound was prepared from the title compound of Example 20 by a standard procedure using trimethylsilylisocyanate in refluxing acetonitrile. The product was obtained as a colorless solid, mp 214-215°C. $[\alpha]_D^{25} = +59.2^\circ$ (c = 1.04, DMF). Analysis calculated for $C_{21}H_{24}N_4O_3 \cdot 0.03 H_2O$: C, 66.20; H, 6.37; N, 14.70. Found: C, 66.10; H, 6.29; N, 14.80.

Example 27

(3R-trans)-N-[2-(N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino)ethyl]methanesulfonamide




The title compound was prepared from the title compound of Example 20 by a standard procedure using methanesulfonyl chloride in pyridine and dichloromethane. The product was obtained as a colorless solid, mp 143-145°C. $[\alpha]_D^{25} = +33.0^\circ$ (c = 1.02, DMF). Analysis calculated for

$C_{21}H_{25}N_3O_4S \cdot 0.67 H_2O$: C, 59.00; H, 6.21; N, 9.83; S, 7.50. Found: C, 59.18; H, 5.80; N, 9.65; S, 7.50.

Example 28

(3R-trans)-N"-Cyano-N-[2-[[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]guanidine



A. (3R-trans)-N'-Cyano-N-[2-[[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]carbamidic acid, phenyl ester

A solution of (3R-trans)-4-[N-(2-aminoethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile (500 mg, 1.48 mmol, the title compound of Example 20) and diphenylcyanocarbonimidate (0.39 g, 1.59 mmol) in acetonitrile (5.0 mL) was heated at reflux temperature for two hours. The reaction mixture was cooled to room temperature and partitioned between ethyl acetate and 1N hydrochloric acid. The organic fraction was separated and the aqueous fraction was reextracted with ethyl acetate. The organic extracts were washed with brine, dried over magnesium sulfate and evaporated to provide the title compound as a colorless gum (0.84 g). The crude material was used for the next reaction.

B. (3R-trans)-N"-Cyano-N-[2-[[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]guanidine


To a solution of the title A compound (0.81 g, 1.47 mmol) in ethanol (9 mL) was added ammonium hydroxide (8.5 mL) and the reaction mixture

was stirred at room temperature for 48 hours. The solvent was evaporated and the residue in ethyl acetate was washed with water and dried over anhydrous magnesium sulfate. The solvent was evaporated and the residue was purified by flash chromatography on silica gel (2% methanol in ethyl acetate). The product was triturated with isopropyl ether to give a colorless amorphous solid, mp 125-130°C (shrinking). $[\alpha]_D^{25} = +31.8^\circ$ (c = 1.12, DMF). Analysis calculated for $C_{22}H_{24}N_6O_2 \cdot 0.2 H_2O \cdot 0.5$ isopropyl ether: C, 63.76; H, 6.33; N, 18.59. Found: C, 63.87; H, 6.35; N, 18.58.

10

Example 29

(3R-trans)-2-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-(2-hydroxyethyl)acetamide



15


The title compound was prepared from (3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid ethyl ester (the title compound of Example 3) and 2-hydroxyethyl amine by the same procedure as described for the title compound of Example 7. The product was obtained as a white foam. $[\alpha]_D^{25} = +32.7^\circ$ (c = 0.62, MeOH). Analysis calculated for $C_{22}H_{25}N_3O_4 \cdot 0.17 H_2O$: C, 66.31; H, 6.41; N, 10.55. Found: C, 66.29; H, 6.42; N, 10.57.

20

25

Example 30

(3R-trans)-4-[4-Chloro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



A. N-(4-Chlorophenyl)-N-[(1H-imidazol-2-yl)methyl]amine

The title compound was prepared from 4-chloroaniline and 2-imidazolecarboxaldehyde by the same procedure as described in Example 24, part


A. The residue was crystallized from ethyl acetate to afford the title compound (1.56 g, 72%) as an off white solid.

B. (3R-trans)-4-[4-Chloro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

The title compound was prepared from (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) and the title A compound by the same procedure as described for the title compound of Example 24. The residue after work up was purified by column chromatography (40% ethyl acetate in hexanes) to afford the title product (198 mg, 28%) as a white solid, mp 266-267°C (softens at 160°C). $[\alpha]_D^{25} = +40.8^\circ$ (c = 0.36, MeOH). Analysis: calculated for $C_{22}H_{21}ClN_4O_3 \cdot 0.23H_2O$: C, 63.9%; H, 5.24; N, 13.56; Cl, 8.58. Found: C, 64.37; H, 5.29; N, 13.17; Cl, 8.24.

Example 31

(3R-trans)-4-[4-Fluoro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, mono hydrochloride



A. N-(4-Fluorophenyl)-N-[(1H-imidazol-2-yl)methyl]amine

The title compound was prepared from 4-fluoroaniline and 2-imidazole-carboxaldehyde by the same procedure as described in Example 24, part

10 A. The product was obtained as a light yellow solid (4.84 g, 95%).

B. (3R-trans)-4-[4-Fluoro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, mono hydrochloride


15 The title compound was prepared from (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) and the title A compound by the same procedure as described for the title compound of Example 24. The colorless product (643 mg, 41%) (mp 252-253 °C, decomposed) in methanol was converted

20 to its hydrochloride by treatment with hydrogen chloride gas in dioxane. The solvent was removed and the residue was dissolved in water (20 mL). The solution was filtered through a Whatman 0.3 μm cellulose nitrate membrane filter and the solvent was removed by freeze drying to afford a white lyophilate. $[\alpha]_D^{25} = -23.3^\circ$ (c = 0.61, MeOH). Analysis calculated for $C_{22}H_{21}FN_3O_2 \cdot HCl \cdot 1.39H_2O$: C, 58.35; H, 5.29; N, 12.37; Cl, 7.83; F, 3.79. Found: C, 58.76; H, 5.12; N, 11.96; Cl, 7.51; F, 4.20.

25

Example 32

(3R-trans)-4-[N-(3-Furanylmethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



A. [N-(3-Furanylmethyl)-N-phenylamine


A mixture of aniline (1.89 g, 19.6 mmol) and 3-furaldehyde (2.50 g, 26.8 mmol) in 1,2-dichloroethane (100 mL) under argon at 5° was treated with sodium triacetoxyborohydride (5.65 g, 26.8 mmol) and acetic acid (1.5 mL). The reaction mixture was stirred at room temperature overnight, concentrated *in vacuo* and the residue was diluted with ethyl acetate. The combined extracts were washed with saturated sodium bicarbonate, dried over MgSO₄ and concentrated. The product was purified by flash chromatography on silical gel (Hexane/EtOAc (20:1) to yield the title compound (3.31 g, 98%). Analysis calculated for C₁₁H₁₁NO: C, 76.28; H, 6.40; N, 8.09. Found: C, 76.58; H, 6.46; N, 8.33.

B. (3R-trans)-4-[N-(3-Furanylmethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

The title compound was prepared from (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) and the title A compound by the same procedure as described for the title compound of Example 1. The product was obtained as a colorless solid (0.85 g, 57%), mp 63-67°C. [α]_D²⁵ = +65.2° (c = 0.71 MeOH). Analysis calculated for C₂₃H₂₂N₂O₄•0.25H₂O: C, 72.90; H, 5.98; N, 7.39. Found: C, 72.94; H, 5.95; N, 7.35.

Example 33

(3R-trans)-4-[N-(2-Furanylmethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



5

A. [N-(2-Furanylmethyl)-N-phenylamine

The title compound was prepared from aniline and 2-furaldehyde by the same procedure as described in Example 32, part A. The product was obtained as an oil (3.43 g, 99%).

10


B. (3R-trans)-4-[N-(2-Furanylmethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

The title compound was prepared from (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) and the title A compound by the same procedure as described for the title compound of Example 1. The product was obtained as a colorless solid (1.07 g, 72%), mp 134-135°C. $[\alpha]_D^{25} = +92.2^\circ$ ($c = 0.78$, MeOH). Analysis calculated for $C_{23}H_{22}N_2O_3$: 0.02 H_2O : C, 73.70; H, 5.93; N, 7.47. Found: C, 73.65; H, 5.63; N, 7.52.

20

Example 34

(3R-trans)-4-[N-[(4,5-Dihydro-2-oxazolyl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile




5

A stirred solution of (3R-trans)-2[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-(2-hydroxyethyl)-acetamide (600 mg, 1.52 mmol, the title compound of Example 29) in dichloromethane (3 mL) at 0°C under argon was treated with triethylamine

- 10 (254 μ L, 1.82 mmol) and methanesulfonyl chloride (130 μ L, 1.67 mmol). The solution was stirred for 30 minutes at -30°C and warmed to room temperature. It was diluted with ethyl acetate and washed with water, sodium bisulfite and brine (100 mL). After drying over anhydrous magnesium sulfate, the solvent was removed. The oily residue in
- 15 dimethylformamide (5 mL) was treated with finely ground potassium carbonate and heated at 150°C for 30 minutes. The reaction mixture was further stirred for two days at room temperature and diluted with ethyl acetate. It was washed with water, 5% sodium bisulfite and brine (100 mL). After drying over anhydrous magnesium sulfate, the solvent
- 20 was removed and the residue was purified by flash chromatography on silica gel (40% ethyl acetate in hexanes). The resulting solid was recrystallized from chloroform-hexanes to give the title product (490 mg, 85%) as fine white needles, mp 218-220°C. $[\alpha]_D^{25} = +71.8^\circ$ (c = 0.4, MeOH). Analysis calculated for $C_{22}H_{23}N_3O_2$: C, 70.00; H, 6.14; N, 11.13. Found: C, 69.96; H, 6.05; N, 11.17.
- 25

Example 35

(3R-(trans)-[(2-Benzoxazolyl)(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4yl)amino]acetic acid, ethyl ester



5

A. (3R-(trans)-4-Amino-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

To a solution of (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (2.5 g, 12.4 mmol, the title A compound of Example 3) in tetrahydrofuran in a re-sealable tube was added concentrated ammonium hydroxide (2 mL). The tube was sealed and the solution was heated on an oil bath at 75°C overnight. The resultant solution was cooled to room temperature, concentrated and extracted with ethyl acetate. The combined organic extracts were dried over anhydrous sodium sulfate and concentrated to yield the title compound as an oil.

15

B. (3R-(trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4yl)amino]acetic acid, ethyl ester

To a solution of (3R-(trans)-4-amino-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile (the title A compound) and ethyl glyoxalate (2.5 g) in methanol (30 mL) and acetic acid (2 mL) at 0°C under argon atmosphere was added sodium cyanoborohydride (1.5 g). The reaction mixture was stirred at 0°C for 30 minutes, poured into saturated sodium bicarbonate (150 mL) and extracted with ethyl acetate. The combined organic extracts were dried over anhydrous sodium sulfate and concentrated. The residue was purified by chromatography to give an oil, which solidified upon standing (1.8 g, 48%).


25

C. (3R-trans)-[(2-Benzoxazolyl)(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4yl)amino]acetic acid, ethyl ester

The title compound was prepared from the title B compound and 2-chlorobenzoxazole by the same procedure as described in Example 19, part B. The product was purified by flash chromatography on silica gel (4:1 mixture of hexane and ethyl acetate) to give the title compound as an amorphous solid (320 mg, 46%), mp -90°C . $[\alpha]_{\text{D}}^{25} = +44.3^{\circ}$ (c = 1, MeOH). Analysis calculated for $\text{C}_{23}\text{H}_{25}\text{N}_3\text{O}_5 \cdot 0.32\text{H}_2\text{O}$: C, 64.6%; H, 5.58 N, 9.83. Found: C, 64.72; H, 5.45; N, 9.77.

Example 36

(3R-trans)-4-[(2-Benzoxazolyl)(2-pyridinylethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



A. (3R-trans)-4-[(2-Pyridinylethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

A solution of (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (1.0 g, 5.0 mmol, the title A compound of Example 3) and 2-aminoethylpyridine (1.1 g, 9.0 mmol) was heated in a sealed tube at 75°C for 18 hours. The reaction mixture was cooled to room temperature and purified by flash chromatography to afford an oil, which solidified upon standing (1.4 g, 87%).

B. (3R-trans)-4-[(2-Benzoxazolyl)(2-pyridinylethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile


The title compound was prepared from the title A compound and 2-chlorobenzoxazole by the same procedure as described in Example 19.

5 part B. The product was purified by flash chromatography on silica gel (4:1 mixture of hexane and ethyl acetate) to give the title compound as a foam (550 mg, 94%). $[\alpha]_D^{25} = +33.2^\circ$ (c = 1, MeOH). Analysis calculated for $C_{26}H_{24}N_4O_3 \cdot 0.93H_2O$: C, 68.29; H, 5.70; N, 12.25. Found: C, 68.64; H, 5.68; N, 11.90.

10

Example 37

(3R-trans)-4-[(2-Benzoxazolyl)(2-furanomethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



15

A. (3R-trans)-4-[(2-Furanomethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

The title compound was prepared from (1aR-cis)-1a,7b-dihydro-2,2-

20 dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) and 2-furfurylamine by the same procedure as described in Example 36, part A. The product was obtained as a colorless oil (1.35 g, 91%).

B. (3R-*trans*)-4-[(2-Benzoxazolyl)(2-furanomethylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile


The title compound was prepared from the title A compound and 2-chlorobenzoxazole by the same procedure as described in Example 19.

part B. The oily product became a foam upon vacuum drying. $[\alpha]_D^{25} = +64.4^\circ$ ($c = 1$, MeOH) Analysis calculated for $C_{24}H_{21}N_3O \cdot 0.40 H_2O \cdot 0.2$ toluene: C, 69.17; H, 5.35; N, 9.53. Found: C, 69.10; H, 5.25; N, 9.43.

10

Example 38

(3R-*trans*)-4-[(2-Furanylmethyl)(2-oxazolylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



15 A. (2,2-Dimethoxyethyl)carbamic acid, 4-nitrophenyl ester

A solution of dimethylaminoacetal (1.1 g, 10 mmol), and methyl amine (1.4 mL, 11 mmol) in a mixture of ether (100 mL) and methylene chloride (5 mL) at 0°C in an ice-bath was treated with 4-nitrophenyl chloroformate (2.2 g, 11 mmol). The solution was slowly warmed up to room temperature and stirred at room temperature for two hours. The reaction mixture was poured into 5% hydrochloric solution (20 mL), the organic layer was separated and dried over anhydrous magnesium sulfate. The solvent was removed to give the title product as a solid, which was used for the next step without further purification.

25

B. (3R-*trans*)-N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl)-2H-1-benzopyran-4-yl)-N'-(2,2-dimethoxyethyl)-N-(2-furanyl)-methylurea

To a solution of (3R-*trans*)-4-[(2-furanomethylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile (1.1 g, 3.69 mmol), the title A compound of Example 37) in acetonitrile (5 mL) was added the

30

title A compound (1.3 g, 4.9 mmol) followed by diisopropylethyl amine
 (0.85 mL). The mixture was heated at reflux temperature over night,
 concentrated in vacuo and the residue was diluted with ethyl acetate
 (100 mL). The resulting solution was washed with 10% aqueous
 5 potassium hydroxide, saturated ammonium chloride and dried over
 magnesium sulfate. The solvent was removed and the residue was
 purified by flash chromatography to provide the title compound as a
 colorless solid. Analysis calculated for $C_{22}H_{27}N_3O_6 \cdot 0.21$ toluene: C,
 62.87; H, 6.45; N, 9.34. Found: C, 62.86; H, 6.58; N, 9.56.

10


**C. [S-(R*,R*)]-N-(2-Furanylmethyl)-N-(2-hydroxy-1-methyl-
 propyl)-N'-(2-oxoethyl)urea**

The solution of the title B compound in acetone (5 mL) was treated with
 10% hydrochloride (1 mL). The mixture was stirred at room temperature
 15 for three hours and neutralized with sodium bicarbonate (3 mL). The
 mixture was concentrated in vacuo and the residue was extracted with
 ethyl acetate (2 x 100 mL). The organic extracts were dried over
 anhydrous magnesium sulfate and concentrated. The residue was purified
 by flash column (3:2/ethyl acetate:hexane) to give an oil (720 mg, 51%
 20 from the title A compound).

**D. (3R-trans)-4-[(2-Furanylmethyl)(2-oxazolyl)amino]-3,4-
 dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile**
 To a solution of the title C compound (700 mg, 1.83 mmol) in methylene
 25 chloride (50 mL) under argon at room temperature was added
 triphenylphosphine (1.0 g, 3.83 mmol), diisopropylethyl amine (1.4 mL,
 7.7 mmol) and iodine (1.0 g, 3.85 mmol). The brown reaction mixture
 was stirred at room temperature for an hour and poured into saturated
 sodiumthiosulfate solution (25 mL). The organic layer was separated,
 30 dried over anhydrous magnesium and the solvent was removed. The
 residue was purified by flash chromatography to give the title compound
 as a colorless foam (260 mg, 39%), mp 63°C. $[\alpha]_D^{25} = +51.7^\circ$ (c 1.0,
 MeOH). Analysis calculated for $C_{20}H_{19}N_3O_4 \cdot 0.15$ ether + 0.4 H₂O: C,
 64.42; H, 5.55; N, 10.94. Found: C, 64.78; H, 5.36; N, 11.15.

Example 39


(3R-*trans*)-4-[N-(Cyanomethyl)phenylamino]-3,4-dihydro-3-hydroxy-
 2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



The title compound was prepared from N-phenylglycinonitrile and (1*aR*-*cis*)-1*a*,7*b*-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) by the procedure described for the title compound of Example 1. The product was titrated with hexanes (containing 1% ethyl acetate) to give a colorless solid, mp 85-90°C. $[\alpha]_D^{25} = +120.9^\circ$ ($c = 0.45$, CDCl_3). Analysis calculated for $\text{C}_{20}\text{H}_{19}\text{N}_3\text{O}_2 \cdot 0.27 \text{ H}_2\text{O} \cdot 0.04 \text{ hexanes}$: C, 71.15; H, 5.93; N, 12.30. Found: C, 71.16; H, 5.81; N, 11.87.

Example 40

(3R-*trans*)-4-[N-(Cyanopropyl)phenylamino]-3,4-dihydro-3-hydroxy-
 2,2-dimethyl-2H-1-benzopyran-6-carbonitrile




The title compound was prepared from anilinopropionitrile and (1*aR*-*cis*)-1*a*,7*b*-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) by the procedure described for the title compound of Example 1. The product was titrated with hexanes (containing 1% ethyl acetate) to give a colorless solid, mp 164-166°C.

$[\alpha]_D = -38.1^\circ$ ($c = 1.0$, CDCl_3). Analysis calculated for $\text{C}_{21}\text{H}_{21}\text{N}_3\text{O}_2 \cdot 0.14 \text{H}_2\text{O}$: C, 72.09; H, 6.13; N, 12.02. Found: C, 71.91; H, 5.83; N, 11.92.

5

Example 41

(3R-trans)-3,4-Dihydro-3-hydroxy-4-[N-(2-methoxyethyl)phenyl-amino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



10 A. N-(2-methoxyethyl)aniline

The title compound was prepared from aniline and methoxyacetaldehyde by the same method as described for Example 5, part A. The residue was purified by flash chromatography on silica gel (ethyl acetate:hexanes/1:15) to give a pale yellow oil.

15


B. (3R-trans)-3,4-Dihydro-3-hydroxy-4-[N-(2-methoxyethyl)-phenyl-amino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

The title compound was prepared from the title A compound and (12R-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno-[c][1]benzopyran-6-carbonitrile (the title A compound of Example 3) by the procedure described for the title compound of Example 1. The oily residue was purified by flash chromatography on silica gel (ethyl acetate:hexanes/1:6), to give a colorless powder (410 mg, 77%), mp 119-124°C. $[\alpha]_D = +54.4^\circ$ ($c = 0.43$, MeOH). Analysis calculated for $\text{C}_{21}\text{H}_{24}\text{N}_2\text{O}_2$: C, 71.57; H, 6.86; N, 7.95. Found: C, 71.53; H, 6.94; N, 7.73.

25

Example 42

(3R-cis)-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-ethyl-acetamide



5

A. (4aR-cis)-1,2,3,4a,5,10b-Hexahydro-5,5-dimethyl-3-oxo-1-phenyl[1]benzopyrano[3,4-b][1,4]oxazine-9-carbonitrile

A solution of N-phenylglycine ethyl ester (900 mg, 5.0 mmol) in 1,2-dichloroethane (10 mL) under argon at 20°C was treated with trimethylaluminum (3.0 mL, 2.0 M in toluene, 6.0 mmol) over 1-2 minutes. After 15 minutes, (1aR-cis)-1a,7b-dihydro-2,2-dimethyl-2H-oxireno[c][1]benzopyran-6-carbonitrile (1.0 g, 5.0 mmol, title A compound of example 3) was added at once and stirring was continued for 1.5 hours.

15 The mixture was diluted with ethyl acetate, quenched with a few drops of water and filtered through celite to remove gelatinous aluminum salts. The filtrate was washed with 5% sodium bicarbonate, water and brine, dried over anhydrous magnesium sulfate and concentrated. The residue was triturated with warm methanol to give warm methanol to give the title product (392 mg), mp 241-243°C. $[\alpha]_D^{25} = -129.1^\circ$ (c = 0.35 CHCl₃). Analysis calculated for: C₃₀H₁₈N₂O₃·0.14 H₂O: C, 71.29; H, 5.47; N, 8.31. Found: C, 71.05; H, 5.30; N, 8.55.

20


B. (3R-cis)-2-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-ethyl-acetamide

A slurry of title A compound (150 mg, 0.45 mmol) in ethanol/chloroform (10 mL, 10:1) was treated with 70% aqueous ethylamine (160 mg, 2.5 mmol), the mixture gradually becoming homogeneous over 0.5 hour. After 2 hours, the solution was concentrated and the residue crystallized from ethyl acetate/hexane (1:10) to give the title product (150 mg, 87%), mp 202-204°C. $[\alpha]_D = +132.3^\circ$ ($c = 0.52$ CHCl₃). Analysis calculated for C₂₂H₂₅N₃O₃: C, 69.64; H, 6.64; N, 11.07. Found: C, 69.32; H, 6.39; N, 10.67.

Compounds of Examples 43 to 106 and 112 to 118 may be prepared by modifying the procedures described in Examples 1 to 42 and 107 to 111 as known by those skilled in the art.

Example 43

(3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(5-methyl-3-isoxazolyl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile




M.P. °C (solvent)	Rotation (c _D) ²⁵	Analysis
80-90 (form)	+36.9 (MeOH)	Calculated for C ₂₁ H ₂₃ N ₃ O ₃ ·0.1 H ₂ O: C, 70.60; H, 5.97; N, 10.74 Found: C, 70.49; H, 5.58; N, 10.43

Example 44

(3R-trans)-4-{4-Fluorophenyl}[(5-methyl-3-isoxazolyl)methyl]-
amino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-
6-carbonitrile

5




M.P. °C (solvent)	Rotation [α] _D ²⁵	Analysis
80-90 (foam)	+12.7 (MeOH)	Calculated for C ₂₃ H ₂₂ FN ₃ O ₃ : C, 67.80; H, 5.44; N, 10.31; F, 4.66. Found: C, 67.84; H, 5.55; N, 10.0%; F, 4.57.

10

Example 45

15 (3S-cis)-2-[N-(6-Cyano-3,4-dihydro-5-hydroxy-2,2-dimethyl-2H-1-
benzopyran-4-yl)phenylamino]-N-ethyl-acetamide




M.P. °C (solvent)	Rotation [α] _D ²⁵	Analysis
204-206 (EtOAc/Hexanes)	-132.5 (CDCl ₃)	Calculated for C ₂₂ H ₂₃ N ₃ O ₃ : C, 69.64; H, 6.64; N, 11.07; Found: C, 69.42; H, 6.14; N, 10.73

20

Example 46


(3R-trans)-[[[5-[[[(3,4-Dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-fluorophenyl)amino]methyl]-2-furanyl]methyl]-amino]acetic acid, ethyl ester, monohydrochloride



M.P. °C (solvent)	Rotation [α] _D ²⁵	Analysis
105-115 (Et ₂ O/Hexanes)	-6.1 (CDCl ₃)	Calculated for C ₂₉ H ₃₀ FN ₃ O ₅ · 1.0 HCl: C, 61.82; H, 5.74; N, 7.72; Cl, 6.52. Found: C, 61.72; H, 5.70; N, 7.47; Cl, 6.09.

Example 47


(3R-trans)-4-[(4-Fluorophenyl)[[5-(hydroxymethyl)-2-furanyl]-methyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation [α] _D ²⁵	Analysis
63-68 (Hexanes)	-50.7 (MeOH)	Calculated for C ₂₇ H ₂₇ FN ₃ O ₄ C, 66.94; H, 5.59; N, 6.51; F, 4.42. Found: C, 67.22; H, 5.81; N, 6.28; Cl, 4.46.

Example 48

(3R-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid, n-butyl ester




M.P. °C (solvent) Rotation [α]_D²⁰ Analysis

82-85 (foam) +80.3 (CDCl₃)

Calculated for C₂₄H₂₈N₂O₄:
C, 70.57; H, 6.91; N, 6.86.
Found: C, 70.50; H, 7.06; N, 6.84.

Example 49

(3R-trans)-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-phenylacetamide




M.P. °C (solvent) Rotation [α]_D²⁰ Analysis

135-145 (Hexanes) -21.7 (MeOH)

Calculated for C₂₆H₂₅N₂O₃:
0.23H₂O = 0.06 hexanes
C, 72.48; H, 6.07; N, 9.62.
Found: C, 72.48; H, 6.03; N, 9.17.

Example 50

(3R-trans)-4-[N-(2-Furanylmethyl)phenylamino]-3,4-dihydro-2,2-dimethyl-6-(1H-tetrazol-5-yl)-2H-1-benzopyran-3-ol




M. P. °C (solvent) Rotation [α]_D²⁵ Analysis

130-135 (Hexanes) +35.9 (MeOH) Calculated for C₂₃H₂₃N₅O₃:
C, 66.17; H, 5.55; N, 16.78.
Found: C, 65.87; H, 5.88;
N, 16.81.

Example 51

(3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(5-methyl-3-isoxazolyl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile




M. P. °C (solvent) Rotation [α]_D²⁵ Analysis

80-90 +36.9 (MeOH) Calculated for C₂₁H₂₁N₃O₃:
C, 70.60; H, 5.97; N, 10.74.
Found: C, 70.49; H, 5.58; N, 10.43.

Example 52

(3S-(trans)-3,4-Dihydro-3-hydroxy-4-[N-[(1H-imidazol-2-yl)methyl]-phenylamino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M. P. °C (solvent) Rotation [α]_D° Analysis


264-265
(decomposition)

-23.9 (MeOH)

Calculated for
C₂₂H₂₂N₄O₂·0.61H₂O:
C, 68.55; H, 6.07; N, 14.54.
Found: C, 68.66; H, 5.99; N, 14.43.

Example 53

(3R-(trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-(2-oxazolylmethyl)phenylamino]-2H-1-benzopyran-6-carbonitrile



M. P. °C (solvent) Rotation [α]_D° Analysis


207-208

+48.8 (CHCl₃)

Calculated for C₂₂H₂₁N₃O₃·
1.00H₂O·0.14EtOAc: C, 68.78;
H, 5.99; N, 10.36.
Found: C, 67.12; H, 5.58; N, 9.92.

Example 54

(3R-trans)-2[[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]methyl]-4-oxazolecarboxylic acid ethyl ester



M.P. °C (solvent) Rotation $[\alpha]_D^{25}$ Analysis


99-100

+16.1 (CHCl₃)

Calculated for C₂₅H₂₅N₃O₅:
C, 67.10; H, 5.63; N, 9.39.
Found: C, 67.21; H, 5.43; N, 9.22.

Example 55

(3R-trans)-2[[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]methyl]-4-oxazolecarboxylic acid mono sodium salt



M.P. °C (solvent) Rotation $[\alpha]_D^{25}$ Analysis


203-204

+33.3 (MeOH)

Calculated for
C₂₃H₂₃NaN₃O₅ (441.40):
C, 60.50; H, 4.79; N, 9.20.
Found: C, 60.67; H, 4.42; N, 9.03.

Example 56

(S*,R*)-N-[[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetyl]-L-serine, methyl ester



M.P. °C (solvent) Rotation [α]_D° Analysis


105-108

+75.6 (CHCl₃)

Calculated for C₂₄H₂₇N₃O₆:
C, 63.55; H, 6.00; N, 9.27.
Found: C, 63.25; H, 6.08; N, 9.15.

Example 57

(3R-trans)-4-[N-[(5-Methyl-1,3,4-oxadiazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent) Rotation [α]_D° Analysis


201-202

+2.1 (CHCl₃)

Calculated for
C₂₂H₂₃N₄O₃·(1/2) 13H₂O:
C, 67.28; H, 5.71; N, 14.27.
Found: C, 67.41; H, 5.80; N, 14.14

Example 58

(3R-trans)-4-[(4-Chlorophenyl)(2-oxazolymethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. 0°C (solvent) Rotation (α)_D²⁰ Analysis


149-150

+06.1 (MeOH)

Calculated for
C₂₂H₂₀ClN₃O₃·0.43H₂O
C, 63.26; H, 5.04; N, 10.06.
Found: C, 63.58; H, 4.71; N, 9.74.

Example 59

(3R-trans)-4-[N-(1H-Benzimidazol-2-ylmethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile




M.P. 0°C (solvent) Rotation (α)_D²⁰ Analysis

246-247 (decomp) -57.7 (CHCl₃)

Calculated for C₂₆H₂₄N₄O₃
C, 73.57; H, 5.70; N, 13.20.
Found: C, 73.52; H, 5.81; N, 12.80

Example 60

(3R-trans)-4-[(2-Benzoxazolyl)(2-(4-morpholinyl)ethyl)amino]-
3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. 0°C (solvent) Rotation [α]_D²⁵ Analysis


110-115

+50.2 (MeOH)

Calculated for C₂₅H₂₉N₄O₄·
1.15 H₂O: C, 63.99; H, 5.51;
N, 11.94.
Found: C, 63.98; H, 6.15; N, 11.92.

Example 61

(3R-trans)-4-[(2-Furanylmethyl)(2-pyrimidinyl)amino]-3,4-dihydro-
3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. 0°C (solvent) Rotation [α]_D²⁵

65-75


+76.8 (MeOH)

Analysis

Calculated for C₂₁H₂₀N₄O₃·
0.85 H₂O: C, 64.39; H, 5.58;
N, 14.30.
Found: C, 64.04; H, 5.17; N, 13.97.

Example 62

(3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[(2-pyrazinyl)-(3-pyridinylmethyl)amino]-2H-1-benzopyran-6-carbonitrile



M. P. °C (solvent)	Rotation [α] _D ²⁵	Analysis
110-112	-9.8 (MeOH)	Calculated for C ₂₂ H ₂₁ N ₅ O ₂ : 0.38H ₂ O-1.42TFA: C, 53.64; H, 4.20; N, 12.59; F, 14.55 Found: C, 53.45; H, 3.79; N, 12.23; F, 14.17.

Example 63


(3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[(3-pyridinylmethyl)(2-pyrimidinyl)amino]-2H-1-benzopyran-6-carbonitrile



M. P. °C (solvent)	Rotation [α] _D ²⁵	Analysis
102-105	+6 (MeOH)	Calculated for C ₂₂ H ₂₁ N ₅ O ₂ : 0.51H ₂ O-1.65TFA: C, 51.98; H, 4.08; N, 11.98; F, 16.09 Found: C, 51.62; H, 3.71; N, 11.78; F, 15.85.

Example 64


(3R-trans)-4-[(2-Benzoxazolyl)(2-pyridinylmethylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation (α) _D °	Analysis
122-124	+49.6 (MeOH)	Calculated for C ₂₅ H ₂₃ N ₄ O ₂ · 0.3C ₆ H ₁₀ O·0.2H ₂ O: C, 69.51; H, 5.61; N, 12.38. Found: C, 69.13; H, 5.27; N, 12.35.

Example 65


(3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[(2-pyrimidinyl)(2-pyridinylmethylamino)-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation (α) _D °	Analysis
227-228	+61.5 (CHCl ₃)	Calculated for C ₂₅ H ₂₃ N ₄ O ₂ · 0.35H ₂ O: C, 67.12; H, 5.55; N, 17.79. Found: C, 67.19; H, 5.25; N, 17.72.

Example 66


(3R-trans)-4-((4-Fluorophenyl)(2-pyridinylmethyl)amino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation (α) _D °	Analysis
182-184	-56.2 (MeOH)	Calculated for C ₂₄ H ₂₂ N ₂ O ₂ F•0.2H ₂ O: C, 70.82; H, 5.55; N, 10.32; F, 4.67. Found: C, 70.73; H, 5.42; N, 10.32; F, 4.52.

Example 67


4-(4-Fluoro-N-1H-imidazol-2-ylmethyl)phenylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation (α) _D °	Analysis
122-124		Calculated for C ₂₂ H ₂₁ N ₃ O ₂ F•0.1C ₆ H ₆ D=0.95H ₂ O: C, 67.10; H, 6.01; N, 13.97; F, 4.75. Found: C, 67.45; H, 6.10; N, 13.55; F, 4.85.

Example 68


(3R-trans)-4-[(4-Fluorophenyl)(2-pyrimidinyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile,
monohydrochloride



M. P. °C (solvent)	Rotation (α) _D ²⁰	Analysis
182-184	-37.2 (CD ₃ OD)	Calculated for C ₂₅ H ₁₉ N ₄ O ₃ F· 0.85 HCl·0.7H ₂ O: C, 60.88; H, 4.93; N, 12.91; Cl, 6.94; F, 4.38. Found: C, 61.25; H, 4.82; N, 12.47; Cl, 6.75; F, 3.93.

Example 69

(3R-trans)-4-[(2-Furanyl)methyl(2-pyrazinyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M. P. °C (solvent)	Rotation (α) _D ²⁰	Analysis
88-90	+100.8 (MeOH)	Calculated for C ₂₁ H ₂₀ N ₄ O ₃ · 0.6 H ₂ O·0.4TFA: C, 60.49; H, 5.03; N, 12.94; F, 5.27. Found: C, 60.51; H, 4.92; N, 12.75; F, 5.41.

Example 70

(3R-trans)-4-[(2-Benzothiazolyl)(3-pyridinylmethylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile




M.P. °C (solvent) Rotation (α)_D° Analysis

134

Calculated for $C_{25}H_{22}N_4O_2S$
 $0.5H_2O + 0.25C_4H_8O$: C, 66.43;
H, 5.47; N, 11.92; S, 6.82.
Found: C, 66.39; H, 5.20; N, 11.72;
S, 6.89.

Example 71

(3R-trans)-4-[(4-Fluorophenyl)(3-pyridinylmethylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent) Rotation (α)_D° Analysis

184-185

-55.0 (MeOH)

Calculated for $C_{24}H_{22}N_2O_2F$
 $0.2CHCl_3$: C, 68.02; H, 5.23;
N, 9.82; F, 4.45.
Found: C, 68.12; H, 5.14; N, 9.48;
F, 4.66.


2132766

HA641a

- 66 -

Example 72

(3R-trans)-4-[(2-Benzothiazolyl)(3-pyridinylmethyl amino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.
1-oxide




M. P. °C (solvent) Rotation (α)_D²⁵ Analysis

195-202

Calculated for C₂₅H₂₂N₄SO₂·
0.76H₂O: C, 63.59; H, 5.02;
N, 11.87; S, 6.79.
Found: C, 63.89; H, 4.86; N, 11.57;
S, 6.22.

Example 73

(3R-trans)-4-[(4-Chlorophenyl)(2-(4-morpholinyl)ethyl amino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.
monohydrochloride



M. P. °C (solvent) Rotation (α)_D²⁵ Analysis

165-170

+30.4 (MeOH)

Calculated for C₂₄H₂₈N₂O₃Cl·
0.84C₄H₁₀O·1.0HCl: C, 60.78;
H, 6.97; N, 7.77; Cl, 13.11.
Found: C, 60.63; H, 6.57; N, 7.32;
Cl, 12.95.


2132766

HA641a

- 67 -

Example 74


(3R-trans)-4-[(4-Fluorophenyl)(2-(4-morpholinyl)ethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile monohydrochloride



M.P. °C (solvent)	Rotation (c/d)	Analysis
170-175	-36.5 (MeOH)	Calculated for $C_{24}H_{25}N_3O_3 \cdot 0.55H_2O$ 0.15 $C_{24}H_{25}N_3O_3 \cdot 1HCl$: C, 60.71; H, 6.57; N, 8.64; F, 3.90; Cl, 8.0; Found: C, 60.75; H, 6.47; N, 8.47; F, 3.59; Cl, 7.94

Example 75


(3R-trans)-4-[(6-Chloro-3-pyridazinyl)(2-(4-morpholinyl)ethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, hydrochloride



M.P. °C (solvent)	Rotation (c/d)	Analysis
>200	-9.5 (MeOH)	Calculated for $C_{22}H_{22}N_4O_3 \cdot Cl$ 1.45 $HCl \cdot 0.29H_2O$: C, 52.69; H, 5.34; N, 13.96; Cl, 17.32; Found: C, 52.93; H, 5.30; N, 13.72; Cl, 17.32

Example 76

(3R-trans)-4-[(2-Benzothiazolyl)(1H-imidazol-2-ylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, monohydrochloride




M.P. °C (solvent) Rotation (α)_D° Analysis

>200 -17.5 (MeOH) Calculated for
C₂₃H₂₁N₃O₂S•1.36HCl•0.56H₂O:
C, 56.24; H, 4.82; N, 14.26;
Cl, 9.82; S, 6.53.
Found: C, 56.62; H, 4.46; N, 13.88;
Cl, 9.83; S, 6.47.

Example 77

(3R-trans)-4-[(6-Chloro-3-pyridazinyl)(1H-imidazol-2-ylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, monohydrochloride




M.P. °C (solvent) Rotation (α)_D° Analysis

210 -1.9 (MeOH) Calculated for
C₂₀H₁₉N₅O₂Cl•HCl•0.64H₂O:
C, 52.36; H, 4.67; N, 18.32;
Cl, 15.46.
Found: C, 52.60; H, 4.39; N, 18.08;
Cl, 15.60.

Example 78

(3R-trans)-4-[(5-Trifluoromethyl-2-pyridinyl)](1H-imidazol-2-ylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, monohydrochloride




5

M.P. °C (solvent)	Rotation (α) _D °	Analysis
180	+2.5 (MeOH)	Calculated for C ₂₂ H ₂₀ N ₃ O ₂ F ₃ ·1.15HCl·0.5C ₄ H ₁₀ O: C, 55.17; H, 5.05; N, 13.41; Cl, 7.80; F, 10.91.
10		Found: C, 55.03; H, 4.27; N, 13.11; Cl, 7.60; F, 10.99.

Example 79


15 (3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[[2-(4-morpholinyl)ethyl](4-phenyl-2-thiazolyl)amino]-2H-1-benzopyran-6-carbonitrile, monohydrochloride



M.P. °C (solvent)	Rotation (α) _D °	Analysis
20 170	-22 (MeOH)	Calculated for C ₂₇ H ₃₀ N ₄ O ₅ ·1.35HCl·H ₂ O: C, 58.13; H, 6.03; N, 10.04; Cl, 8.58; S, 5.75.
25		Found: C, 58.38; H, 5.37; N, 9.60; Cl, 8.68; S, 6.19.

Example 80

(3R-trans)-3,4-Dihydro-3-hydroxy-4-((1H-imidazol-2-yl-methyl)
(4-phenyl-2-thiazolyl)amino)-2,2-dimethyl-2H-1-benzopyran-6-
carbonitrile




5

M. P. °C (solvent)	Rotation [α] _D ²⁵	Analysis
160-170	+16.0 (MeOH)	Calculated for C ₂₅ H ₂₃ N ₃ O ₂ S·0.2H ₂ O: 0.1C ₄ H ₈ O ₂ : C, 64.91; H, 5.19, N, 14.91; S, 6.82. Found: C, 65.23; H, 5.14; N, 14.49, S, 7.61.

10

Example 81

15 (3R-trans)-3,4-Dihydro-3-hydroxy-4-((1H-imidazol-2-yl-methyl)-
(4-methyl-2-thiazolyl)amino)-2,2-dimethyl-2H-1-benzopyran-6-
carbonitrile



20


M. P. °C (solvent)	Rotation [α] _D ²⁵	Analysis
178-183	+7.4 (MeOH)	Calculated for C ₂₆ H ₂₁ N ₃ O ₂ S·0.15H ₂ O: 0.17C ₄ H ₈ O ₂ : C, 60.12; H, 5.53; N, 16.95; S, 7.76. Found: C, 60.10; H, 5.37; N, 16.59; S, 7.75.

25

Example 82

(3R-trans)-N-[2-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]-2,2-dimethyl-propionamide

5




M.P. °C (solvent)	Rotation (α) _D °	Analysis
194-195 (hexanes)	+61.1 (DMF)	Calculated for C ₂₅ H ₃₁ N ₃ O ₂ ·0.15H ₂ O: C, 70.78; H, 7.44; N, 9.91. Found: C, 70.89; H, 7.42; N, 9.80.

10

Example 83


15 (3R-trans)-N-[2-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]-N'-phenylurea



M.P. °C (solvent)	Rotation (α) _D °	Analysis
20 193-195	+32.3 (DMF)	Calculated for C ₂₇ H ₂₆ N ₄ O ₂ ·0.34H ₂ O: C, 70.10; H, 6.25; N, 12.11. Found: C, 69.72; H, 6.07; N, 11.79.

Example 84


(3R-trans)-N-[2-[N-(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]-1-pyrrolidinecarboxamide



M. P. °C (solvent)	Rotation $[\alpha]_D^{25}$	Analysis
125-135	+35.8 (DMF)	Calculated for $C_{25}H_{30}N_4O_3 \cdot 0.38H_2O$: C, 68.03; H, 7.02; N, 12.69. Found: C, 68.13; H, 7.05; N, 12.21.

Example 85


(3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(2-oxo-1-pyrrolidinyl)ethyl]phenyl]amino]-2H-1-benzopyran-6-carbonitrile



M. P. °C (solvent)	Rotation $[\alpha]_D^{25}$	Analysis
214-216	+14.4 (DMF)	Calculated for $C_{24}H_{27}N_3O_3 \cdot 0.18H_2O$: C, 70.53; H, 6.75; N, 10.28. Found: C, 70.57; H, 6.57; N, 10.24.

Example 86


(3R-trans)-[[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-fluorophenyl)amino]methyl]phosphonic acid, diethyl ester



M.P. °C (solvent)	Rotation [α] _D °	Analysis
55-56	+7.2 (MeOH)	Calculated for C ₂₃ H ₂₄ N ₂ FPO ₅ •0.27H ₂ O: C, 59.12; H, 6.16; N, 5.99; F, 4.07; P, 6.63. Found: C, 59.23; H, 6.15; N, 5.88; F, 3.86; P, 6.43.

Example 87


[N-(4-Chlorophenyl)-N-(6-cyano-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-4-yl)amino]acetic acid, ethyl ester



M.P. °C (solvent)	Rotation [α] _D °	Analysis
55-57		Calculated for C ₂₂ H ₂₁ N ₂ ClO ₄ • 0.02CHCl ₃ : C, 65.32; H, 5.83; N, 6.92; Cl, 9.28. Found: C, 65.38; H, 5.79; N, 6.90; Cl, 9.49.

Example 88

4-[(4-Chlorophenyl)(1H-imidazol-2-ylmethylamino)-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent) Rotation [α]_D²⁰ Analysis


212-214

Calculated for C₂₂H₂₁N₄ClO:
0.42H₂O: C, 66.16; H, 5.26;
N, 14.03.

Found: C, 66.42; H, 5.33; N, 13.77.

Example 89

(3R-(trans))-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(3-methyl-1,2,4-oxadiazol-5-yl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent) Rotation [α]_D²⁰ Analysis

148-149


+60.6 (MeOH)

Calculated for C₂₂H₂₂N₄O₃:
C, 67.68; H, 5.68; N, 14.35.

Found: 67.74; H, 5.58; N, 14.34.

Example 90

(3R-trans)-4-[(4-Chlorophenyl)((3-methyl-1,2,4-oxadiazol-5-yl)-methylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent) Rotation [α]_D° Analysis

102-105 (hexanes) +79 (MeOH)

Calculated for C₂₂H₂₁N₄ClO₃·

0.24 H₂O: C, 61.57; H, 5.04;


N, 13.05; Cl, 8.26.

Found: C, 62.00; H, 5.09; N, 12.62.

Cl, 8.64

Example 91

(3R-trans)-4-[(4-Fluorophenyl)((3-methyl-1,2,4-oxadiazol-5-yl)-methylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent) Rotation [α]_D° Analysis

116-118 (hexanes) +37.6 (MeOH)

Calculated for C₂₂H₂₁N₄FO₃·

0.02 H₂O: C, 64.65; H, 5.19;

N, 13.71; F, 4.65.


Found: C, 64.65; H, 5.19; N, 13.71.

F, 4.65.

Example 92

(3R-trans)-[N-(3,4-Dihydro-3-hydroxy-2,2-dimethyl-7-(trifluoromethyl)-2H-1-benzopyran-4-yl)]phenylamino]acetic acid, ethyl ester

5




M.P. °C (solvent)	Rotation (α) _D °	Analysis
100-103	-102 (CHCl ₃)	Calculated for C ₂₂ H ₂₄ NF ₃ O ₄ : C, 62.19; H, 5.73; N, 3.30. Found: C, 62.33; H, 5.54; N, 3.16

10

Example 93

(3R-trans)-3,4-Dihydro-3-hydroxy-4-[N-[[3-(hydroxymethyl)-1,2,4-oxadiazol-5-yl]methyl]phenylamino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

15




M.P. °C (solvent)	Rotation (α) _D °	Analysis
95-98 (hexanes)	+69.8 (MeOH)	Calculated for C ₂₂ H ₂₂ N ₄ O ₄ : C, 65.01; H, 5.46; N, 13.78. Found: C, 64.77; H, 5.46; N, 13.46

20

Example 94

(3R-trans)-[N-(3,4-Dihydro-3-hydroxy-2,2-dimethyl-8-(trifluoromethyl)-2H-1-benzopyran-4-yl)]phenylamino]acetic acid, ethyl ester

5




M. P. °C (solvent)	Rotation (α) _D ²⁰	Analysis
45-48	+82 (CHCl ₃)	Calculated for C ₂₂ H ₂₄ NF ₃ O ₄ · 0.17 H ₂ O: C, 61.96; H, 5.75; N, 3.28 Found: C, 62.34; H, 5.52; N, 2.90

10

Example 95

(3R-trans)-4-[(4-Chlorophenyl)[(3-(hydroxymethyl)-1,2,4-oxadiazol-5-yl)methyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile

15




M. P. °C (solvent)	Rotation (α) _D ²⁰	Analysis
101-103 (hexanes)	+83 (MeOH)	Calculated for C ₂₇ H ₂₁ N ₄ ClO ₄ · 0.19 H ₂ O: C, 59.46; H, 4.85; N, 12.61; Cl, 7.98 Found: C, 59.82; H, 4.77; N, 12.25; Cl, 7.58

20

25

Example 96


(3R-trans)-4-[(4-Fluorophenyl)[(3-(hydroxymethyl)-1,2,4-oxadiazol-5-yl)methyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M. P. °C (solvent)	Rotation $[\alpha]_D^{25}$	Analysis
115-118	+40.3 (MeOH)	Calculated for $C_{22}H_{21}FN_4O_4 \cdot 0.75 H_2O$: C, 60.35; H, 5.18; N, 12.79; F, 4.34. Found: C, 60.73; H, 4.85; N, 12.41; F, 4.09.

Example 97


(3R-trans)-4-[N-[(3-Amino-1,2,4-oxadiazol-5-yl)methyl]-phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M. P. °C (solvent)	Rotation $[\alpha]_D^{25}$	Analysis
203-205	+51.4 (MeOH)	Calculated for $C_{21}H_{21}N_5O_4 \cdot 0.5 H_2O$: C, 62.99; H, 5.41; N, 17.49. Found: C, 63.43; H, 5.18; N, 17.05.

Example 98


(3R-trans)-[N-(6-Benzoyl-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid, ethyl ester



M.P. °C (solvent)	Rotation (α) _D ²⁰	Analysis
144-145	+58.9 (MeOH)	Calculated for C ₂₄ H ₂₆ NO ₃ ·0.19 H ₂ O: C, 72.65; H, 6.40; N, 3.03.
		Found: C, 72.72; H, 6.70; N, 2.96.

Example 99

(3R-trans)-[(6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-fluoro-3-[(phenylmethoxy)carbonyl]phenyl)amino]acetic acid, ethyl ester




M.P. °C (solvent)	Rotation (α) _D ²⁰	Analysis
61-61	+31.8 (MeOH)	Calculated for C ₃₀ H ₂₆ FN ₂ O ₅ ·1.24 H ₂ O: C, 64.93; H, 5.72; N, 5.05.
		Found: C, 65.24; H, 5.33; N, 4.74.

Example 100

(3R-trans)-[5-[[[6-Cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl](4-fluorophenylamino)methyl]-1,2,4-oxadiazol-3-yl]methoxy]acetic acid

5



M.P. °C (solvent) Rotation (α)_D²⁵ Analysis

125-128

-30 (MeOH)

Calculated for C₂₄H₂₃FN₄O₆1.47 H₂O: C, 58.72; H, 4.91;

N, 11.41; F, 3.87.

Found: C, 59.15; H, 4.22; N, 10.98;


F, 3.55

10

Example 101

(3R-trans)-[5-[[[4-Chlorophenyl](6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)amino]methyl]-1,2,4-oxadiazol-3-yl]methoxy]acetic acid

15



M.P. °C (solvent) Rotation (α)_D²⁵ Analysis

145-148

-51.7 (MeOH)

Calculated for C₂₄H₂₁ClN₄O₆

C, 57.78; H, 4.65; N, 11.23;

Cl, 7.11.


Found: C, 57.45; H, 4.74; N, 10.98;

Cl, 7.07

25

Example 102


(3R-trans)-[(3-Carboxy-4-fluorophenyl)(6-hydroxy-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)amino]acetic acid, ethyl ester



M. P. °C (solvent)	Rotation (c/d)	Analysis
195-197	-51.4 (MeOH)	Calculated for $C_{21}H_{23}FN_2O_6 \cdot 0.35 H_2O$: C, 59.40; H, 5.06; N, 5.93 Found: C, 59.70; H, 4.76; N, 5.60

Example 103

(3R-trans)-4-[(4-Fluorophenyl)((2-methyl-2H-tetrazol-5-yl)methyl)-amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile




M. P. (°C) (solvent)	Rotation (c/d)	Analysis
20 145-148		Calculated for $C_{21}H_{21}FN_5O_2$: C, 61.76; H, 5.18; N, 20.58 Found: C, 61.87; H, 4.96; N, 20.74

Example 104

(3R-trans)-4-[(4-Fluorophenyl)(1-methyl-1H-tetrazol-5-yl)methyl]-
amino-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-
carbonitrile

5



M.P. °C (solvent) Rotation (c/d)° Analysis

130-135

Calculated for $C_{21}H_{21}FN_4O_2$:

C, 61.76; H, 5.18; N, 20.58.


Found: C, 61.62; H, 5.23; N, 19.85

10

Example 105

(3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-(2-
methylsulfinylethyl)phenylamino]-2H-1-benzopyran-6-carbonitrile

15



M.P. °C (solvent) Rotation (c/d)° Analysis

75-85 (shrnks)

-122 (CHCl₃)Calculated for $C_{21}H_{24}N_2O_2S$ +
0.46 H₂O: C, 64.21; H, 6.40;

N, 7.13; S, 8.16


Found: C, 64.26; H, 6.38; N, 7.08;

S, 7.88

20

Example 106


(3R-(trans))-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-(2-(methylsulfonyl)ethyl)phenylamino]-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation [α] _D ²⁵	Analysis
75-85 (shrinks)	-76.9 (CHCl ₃)	Calculated for C ₂₁ H ₂₄ N ₂ O ₄ S• 0.58 H ₂ O: C, 61.37; H, 6.17; N, 6.82; S, 7.80. Found: C, 61.76; H, 6.15; N, 6.43; S, 7.53.

Example 107

(3R-(trans))-4-[4-Chloro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile monohydrochloride



A. N-(4-Chlorophenyl)-N-[(1H-imidazol-2-yl)methyl]amine

- 20 A mixture of 4-chloroaniline (66.55 g, 522.43 mmol) and 2-imidazolecarboxaldehyde (50.2 g, 522.43 mmol) in methanol (1000 ml.) was stirred at 55-60°C overnight. The light brown reaction mixture was cooled in an ice bath and treated with sodium borohydride (21.74 g, 574.67 mmol) in small portions. The reaction mixture was allowed to
- 25 warm to room temperature and stirred for two hours. It was concentrated and partitioned between water (~500 ml.) and ethyl acetate (1200 ml.), giving a white solid/aqueous layer and a brown organic layer. The organic layer was removed and the aqueous mixture was reextracted with ethyl

acetate (3x200 mL). The combined organic layers were washed with brine, dried over sodium sulfate and concentrated. The resulting mixture was treated with hexanes and stored in the freezer for two hours. The white solid was collected by filtration and washed with cold ethyl acetate/hexane (2:1) to provide the title product (83.36 g, 77%) as a white solid, mp 163-165°C. Analysis calculated for $C_{10}H_{10}ClN_3$: C, 57.84; H, 4.85; N, 20.23; Cl, 17.07. Found: C, 57.82; H, 4.85; N, 20.04; Cl, 16.77.

10 **B. (3R-trans)-4-[4-Chloro-N-[(1H-imidazol-2-yl)methyl]-phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile**


A mixture of title A compound (41.3 g, 198.79 mmol), title A compound of Example 3 (40.0 g, 198.79 mmol) and anhydrous cobalt chloride (25.8 g, 198.79 mmol) in dry acetonitrile (160 mL) under argon was heated at 60°C (oil bath temperature) for 28 hours. The reaction mixture was allowed to cool to room temperature and treated with saturated sodium bicarbonate (600 mL) followed by ethyl acetate (1600 mL). The well shaken mixture was filtered through a short pad of celite, yellow organic layer was separated and washed with brine. After drying over anhydrous sodium sulfate, the solvent was removed and the residue was treated with hexanes (1000 mL) and ethyl acetate (100 mL). The mixture was heated on a steam bath (10-20 minutes), allowed to cool to room temperature and filtered to afford a white solid. This material was heated with methanol (2000 mL) for 15 minutes, allowed to cool to room temperature and filtered to provide (3R-trans)-4-[4-Chloro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile (44.72 g, 55%) as a colorless solid mp 266-267°C. $[\alpha]_D^{25} = +46.9^\circ$ (c = 1.15, acetone).

C. (3R-trans)-4-[4-Chloro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, mono hydrochloride

- A solution of title B compound (50.0 g, 122.29 mmol) in anhydrous tetrahydrofuran (800 mL) at 0°C was treated with a freshly prepared 4.0N hydrogen chloride gas in diethyl ether (36.6 mL, 178.61 mmol). The solution was stirred at 0°C for ten minutes and the solvent was removed to give a yellow oil which was treated with diethyl ether to give the title product (55.87 g, 98%) as a white solid, mp 189-190°C. Analysis calculated for $C_{22}H_{21}ClN_4O_2 \cdot HCl \cdot 0.40H_2O \cdot 0.20THF$: C, 58.64; H, 5.27; N, 12.00; Cl, 15.18. Found: C, 58.72; H, 5.39; N, 11.72; Cl, 15.46. $[\alpha]_D^{25} = +9.5^\circ$ (c = 1.00, MeOH)

Example 108


- 15 (3R-trans)-4-[4-Chloro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, hydrogensulfate



- 20 The title compound was prepared from the title B compound of Example 107 by treatment with concentrated sulfuric acid in tetrahydrofuran. The solvent was removed and the product was triturated with ether to give a colorless solid, mp 154-156°C. Analysis calculated for $C_{22}H_{21}ClN_4O_2 \cdot H_2SO_4 \cdot 0.9H_2O \cdot 0.30Et_2O$: C, 51.09; H, 5.14; N, 10.27; Cl, 6.50; S, 5.88. Found: C, 50.89; H, 4.83; N, 10.01; Cl, 6.67; S, 6.17. $[\alpha]_D^{25} = +6.8^\circ$ (c = 1.12, MeOH)

Example 109


(3R-trans)-4-[4-Chloro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, methanesulfonate



- 5 The title compound was prepared from the title B compound of Example 107 by treatment with methanesulfonic acid in tetrahydrofuran. The solvent was removed and the product was triturated with ether to give a colorless solid, mp 155-156°C. Analysis calculated for
- 10 $C_{22}H_{21}ClN_4O_2 \cdot MeSO_3H \cdot 1.0H_2O \cdot 0.06Et_2O$: C, 52.92; H, 5.27; N, 10.62; Cl, 6.72; S, 6.08. Found: C, 52.92; H, 5.17; N, 10.23; Cl, 6.50; S, 6.57. $[\alpha]_D^{25} = +8.0^\circ$ (c = 1.31, MeOH)

Example 110


- 15 (3R-trans)-4-[4-Chloro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, phosphate



- 20 The title compound was prepared from the title B compound of Example 107 by treatment with phosphoric acid in tetrahydrofuran. The solvent was removed and the product was triturated with ether to give a colorless solid, mp 160-161°C. Analysis calculated for
- 25 $C_{22}H_{21}ClN_4O_2 \cdot H_3PO_4 \cdot 3.4H_2O \cdot 0.35Et_2O$: C, 47.31; H, 5.82; N, 9.43; Cl, 5.87. Found: C, 47.34; H, 5.06; N, 9.07; Cl, 5.92. $[\alpha]_D^{25} = +18.8^\circ$ (c = 1.18, MeOH).

Example 111


(3R-trans)-4-[4-Chloro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, nitrate



The title compound was prepared from the title B compound of Example 107 by treatment with nitric acid in tetrahydrofuran. The solvent was removed and the product was triturated with ether to give a colorless solid, mp 141-142°C (decomposition). Analysis calculated for $C_{22}H_{21}ClN_4O_2 \cdot HNO_3 \cdot 0.48H_2O \cdot 0.20Et_2O$: C, 55.28; H, 5.08; N, 14.14; Cl, 7.16. Found: C, 55.29; H, 5.11; N, 13.82; Cl, 7.05. $[\alpha]_D^{25} = +8.3$ ($c = 0.99$, MeOH).

Example 112

[3R-(3 α ,4 β (R*))]-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-[[5-oxo-1-(phenylmethyl)-2-pyrrolidinyl]methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile



M.P. (solvent) Rotation (mp) Analysis

161-163


-102 (CHCl₃)

Calculated for $C_{24}H_{23}N_3O_2$
0.53 H₂O: C, 73.36; H, 6.58;
N, 8.55

Found: C, 73.59; H, 6.49; N, 8.32

Example 113


[3R-[3 α ,4 β (S*)]]-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(5-oxo-2-pyrrolidinyl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation [α] _D °	Analysis
186-189	-116 (CHCl ₃)	Calculated for C ₂₃ H ₂₅ N ₃ O ₃ · 0.50 H ₂ O: C, 68.98; H, 6.54; N, 10.49. Found: C, 69.16; H, 6.49; N, 10.31

Example 114


[3R-[3 α ,4 β (R*)]]-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(5-oxo-2-pyrrolidinyl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation [α] _D °	Analysis
amorphous solid	-34 (CHCl ₃)	Calculated for C ₂₃ H ₂₅ N ₃ O ₃ · 0.86 H ₂ O: C, 68.04; H, 7.04; N, 9.72. Found: C, 68.03; H, 6.68; N, 9.68

Example 115


(3R-trans)-4-[N-[(1,5-Dimethyl-1H-pyrazol-3-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation (α) _D ²⁵	Analysis
194-199	+27.1 (MeOH)	Calculated for C ₂₄ H ₂₆ N ₄ O ₂ •0.23 H ₂ O: C, 70.90; H, 6.56; N, 13.78. Found: C, 70.99; H, 6.55; N, 13.69

Example 116


(3R-trans)-3,4-Dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(5-methyl-1H-pyrazol-3-yl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation (α) _D ²⁵	Analysis
110-120	+30.8 (MeOH)	Calculated for C ₂₄ H ₂₄ N ₄ O ₂ •0.78 H ₂ O: C, 68.63; H, 6.40; N, 13.92. Found: C, 69.06; H, 6.46; N, 13.50

Example 117


(3R-trans)-4-[N-[(1,3-Dimethyl-1H-pyrazol-5-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile



M.P. °C (solvent)	Rotation (alq)	Analysis
108-115	+7.9 (MeOH)	Calculated for $C_{24}H_{26}N_4O_2 \cdot 0.50$ H ₂ O-0.30 $C_{48}H_{52}N_8O_4$ C, 69.11; H, 6.77; N, 12.79 Found: C, 69.20; H, 6.63; N, 12.74

Example 118

[3R-[3 α ,4B-(Z)]]-4-[N-(2-Amino-4-oxo-2-pentenyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile




M.P. °C (solvent)	Rotation (alq)	Analysis
188-190	52.0 (MeOH)	Calculated for $C_{21}H_{22}N_2O_4$ C, 70.57; H, 6.44; N, 10.73 Found: C, 70.34; H, 6.52; N, 10.47

What we claim is:

1. A compound of the formula

I




or pharmaceutically acceptable salts thereof wherein

a, b and d are all carbon atoms or one of a, b and d is a nitrogen atom or -N(O)- and the others are carbon atoms.



Y is a single bond, -CH₂-, -C(O)-, -O-, -S- or -NR⁶-.

R¹ is aryl or heterocyclo.

R² is -COOR⁷, -CO-amino, -CO-substituted amino, amino, substituted amino, -NR⁸CO-amino, -NR⁸CO-substituted amino, -NR⁸COR⁹, -NR⁸SO₂R⁹, -NR⁸(C=NCN)-amino, -NR⁸(C=NCN)-substituted amino,



-SR³, -SOR³, -SO₂R³, -OR³, cyano, heterocyclo, pyridine N-oxide,





R³ is hydrogen, hydroxy or -OK(O)R³.

R⁴ and R⁵ are each independently hydrogen, alkyl or arylalkyl or R⁴ and R⁵ taken together with the carbon atom to which they are attached form a 5- to 7-membered carbocyclic ring.

R⁶ is hydrogen, alkyl, haloalkyl, alkenyl, alkynyl,

cycloalkyl, arylalkyl, cycloalkylalkyl, -CN, -NO₂, -COR⁷, -COOR⁸,





$-\text{OCH}_2\text{CF}_3, -\text{OCOalkyl}, -\text{OCONR}^8\text{alkyl}, -\text{NR}^8\text{COalkyl}, -\text{NR}^8\text{COOalkyl}$
or $-\text{NR}^8\text{CONR}^9$, tetrazolyl, imidazolyl, oxazole or triazole;

R^7 is hydrogen, alkyl, hydroxy, $-\text{O-alkyl}$, amino,
substituted amino, $-\text{NHCOR}^8$, $-\text{CN}$ or $-\text{NO}_2$;

R^8 and R^9 are independently hydrogen, alkyl, haloalkyl,
aryl, arylalkyl, cycloalkyl or (cycloalkyl)alkyl;

X is alkyl or X-R^2 together can be hydrogen, aryl or
heterocyclo when R^1 is heterocyclo, and

n is an integer of 1 to 3.

2. The compounds as recited in Claim 1 wherein

a, b and d are carbon atoms.

X is alkyl.

Y is a single bond or $-\text{O}-$.

R^1 is aryl or heterocyclo;

R^2 is $-\text{COOR}^3$, $-\text{COamino}$, $-\text{CO-substituted amino}$,

$-\text{NHCOCH}_3$, $-\text{NHCO}_2\text{Me}$, $-\text{NHCONH}_2$, $-\text{NH}_2\text{C}=\text{NCN}(\text{NH}_2)$, imidazole,
furan, pyridine, oxazole, hydroxy, $-\text{NHCO-substituted amino}$ or $-\text{SO}_2\text{Me}$.

R^3 is hydroxy;

R^4 and R^5 are methyl;

R^6 is cyano, $-\text{NO}_2$, $-\text{CF}_3$, halo, alkyl or tetrazol, and

R^7 is hydrogen

3. The compounds as recited in Claim 1, which are

trans-[6-cyano-3,4-dihydro-3'-hydroxy-2,2-dimethyl-2H-1-benzopyran-4'-ylphenylamino]acetic acid, ethyl ester;

(1S-trans)-[6-cyano-3,4-dihydro-3'-hydroxy-2,2-dimethyl-2H-1-benzopyran-4'-ylphenylamino]acetic acid, ethyl ester;

(1R-trans)-[6-cyano-3,4-dihydro-3'-hydroxy-2,2-dimethyl-2H-1-benzopyran-4'-ylphenylamino]acetic acid, ethyl ester;

trans-[6-cyano-3,4-dihydro-3'-hydroxy-2,2-dimethyl-2H-1-benzopyran-4'-ylphenylamino]acetic acid

(3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-fluorophenyl)amino]acetic acid, ethyl ester.

(3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-chlorophenyl)amino]acetic acid, ethyl ester.

(3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetamide.

(3S-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[(4-phenyl-2-thiazolyl)amino]-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[(4-phenyl-2-thiazolyl)amino]-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-[N-(3,4-dihydro-3-hydroxy-2,2-dimethyl-6-(1H-tetrazol-5-yl)-2H-1-benzopyran-4-yl)phenylamino]acetic acid, ethyl ester.

(3R-trans)-2-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-ethylacetamide.

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N(2-(1-pyrroloindinyl)-2-oxoethyl)phenylamino]-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N(2-(4-morpholinyl)-2-oxoethyl)phenylamino]-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-(2-furanylmethyl)acetamide.

(3R-trans)-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-(2-(4-morpholinyl)ethyl)acetamide.

(3R-trans)-4-[(4-fluorophenyl)(2-hydroxy-2-methylpropyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-[3a,4b R*])-4-[(4-fluorophenyl)(2-hydroxypropyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-methyl-2-thiazolyl)amino]acetic acid, ethyl ester.

(3R-trans)-4-[N-(2-benzoxazolyl)-N-(2,2-dimethoxyethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-4-[N-(1,3-dihydro-1,3-dioxo-2H-isindol-2-yl)ethyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-4-[N-(2-aminoethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]butyric acid, ethyl ester.

(3R-trans)-3-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]propanoic acid, ethyl ester.

(3R-trans)-3,4-dihydro-3-hydroxy-4-[N-[(1H-imidazol-2-yl)methyl]phenylamino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-4-[[2-(acetylaminoethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-[2-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]urea.

(3R-trans)-N-[2-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]methanesulfonamide.

(3R-trans)-N'-cyano-N-[2-[[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]guanidine.

(3R-trans)-2-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-(2-hydroxyethyl)acetamide.

(3R-trans)-4-[4-chloro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-4-[4-fluoro-N-[(1H-imidazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, mono hydrochloride.

(3R-trans)-4-[N-(2-furanylmethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-4-[N-(2-furanylmethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-4-[N-[(4,5-dihydro-2-oxazol-1-en-5-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-[(2-benzoxazolyl)-6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl]amino]acetic acid, ethyl ester.

(3R-trans)-4-[(2-benzoxazolyl)-2-pyridylethyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile.

(3R-trans)-4-[(2-benzoxazolyl)(2-furanomethylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(2-furanylmethyl)(2-oxazolylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[N-(cyanomethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[N-(cyanoethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-3,4-dihydro-3-hydroxy-4-[N-(2-methoxyethyl)phenylamino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-cis)-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-ethylacetamide;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-(5-methyl-3-isoazolylmethyl)phenylamino]-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(4-fluorophenyl)(5-methyl-3-isoazolylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3S-cis)-2-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-ethylacetamide;

(3R-trans)-{[(5-[(3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-fluorophenylamino)methyl]-2-furanylmethyl)amino]acetic acid, ethyl ester, monohydrochloride};

(3R-trans)-4-[(4-fluorophenyl)(5-hydroxymethyl)-2-furanylmethylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-[6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid, n-butyl ester;

(3R-trans)-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]-N-phenylacetamide;

(3S-trans)-4-[N-(2-furanylmethyl)phenylamino]-3,4-dihydro-2,2-dimethyl-6-1H-tetrazol-5-yl-2H-1-benzopyran-3-yl;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-(5-methyl-3-isoazolylmethyl)phenylamino]-2H-1-benzopyran-6-carbonitrile;

(3S-trans)-3,4-dihydro-3-hydroxy-4-[N-[(1H-imidazol-2-yl)methyl]phenylamino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-(2-oxazolylmethyl)phenylamino]-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-2[[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]methyl]-4-oxazolecarboxylic acidethyl ester;

(3R-trans)-2[[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]methyl]-4-oxazolecarboxylic acid mono sodium salt;

(S*,R*)-N-[[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetyl]-L-serine, methyl ester;

(3R-trans)-4-[N-[(5-methyl-1,3,4-oxadiazol-2-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(4-chlorophenyl)(2-oxazolylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[N-(1H-benzimidazol-2-ylmethyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(2-benzoxazolyl)(2-(4-morpholinyl)ethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(2-furanylmethyl)(2-pyrimidinyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[(2-pyrazinyl)-(3-pyridinylmethyl)amino]-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[(3-pyridinylmethyl)(2-pyrimidinyl)amino]-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(2-benzoxazolyl)(2-pyridinylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[(2-pyrimidinyl)(2-pyridinylmethyl)amino]-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(4-fluorophenyl)(2-pyridinylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

4-[4-fluoro-N-(1H-imidazol-2-ylmethyl)phenylamino]-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(4-fluorophenyl)(2-pyrimidinyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, monohydrochloride;

(3R-trans)-4-[(2-furanylmethyl)(2-pyrazinyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(2-benzothiazolyl)(3-pyridinylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(4-fluorophenyl)(3-pyridinylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(2-benzothiazolyl)(3-pyridinylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, 1-oxide;

(3R-trans)-4-[(4-chlorophenyl)[2-(4-morpholinyl)ethyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, monohydrochloride;

(3R-trans)-4-[(4-fluorophenyl)[2-(4-morpholinyl)ethyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, monohydrochloride;

(3R-trans)-4-[(6-chloro-3-pyridazinyl)[2-(4-morpholinyl)ethyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, hydrochloride;

(3R-trans)-4-[(2-benzothiazolyl)(1H-imidazol-2-ylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, monohydrochloride;

(3R-trans)-4-[(6-chloro-3-pyridazinyl)(1H-imidazol-2-ylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, monohydrochloride;

(3R-trans)-4-[(5-trifluoromethyl-2-pyridinyl)(1H-imidazol-2-ylmethyl)amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, monohydrochloride;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[[2-(4-morpholinyl)ethyl](4-phenyl-2-thiazolyl)amino]-2H-1-benzopyran-6-carbonitrile, monohydrochloride;

(3R-trans)-3,4-dihydro-3-hydroxy-4-[(1H-imidazol-2-yl-methyl)(4-phenyl-2-thiazolyl)amino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-3,4-dihydro-3-hydroxy-4-[(1H-imidazol-2-yl-methyl)-(4-methyl-2-thiazolyl)amino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-N-[2-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]-2,2-dimethyl-propanamide;

(3R-trans)-N-[2-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]-N'-phenylurea;

(3R-trans)-N-[2-[N-(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]ethyl]-1-pyrrolidine-carboxamide;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(2-oxo-1-pyrrolidinyl)ethyl]phenyl]amino]-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-[[[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-fluorophenyl)amino]methyl]phosphonic acid, diethyl ester;

[N-(4-chlorophenyl)-N-(6-cyano-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-4-yl)amino]acetic acid, ethyl ester;

4-[(4-chlorophenyl)(1H-imidazol-2-ylmethyl)amino]-3,4-dihydro-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(3-methyl-1,2,4-oxadiazol-5-yl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(4-chlorophenyl)(3-methyl-1,2,4-oxadiazol-5-yl)methyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(4-fluorophenyl)(3-methyl-1,2,4-oxadiazol-5-yl)methyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-[N-[3,4-dihydro-3-hydroxy-2,2-dimethyl-7-(trifluoromethyl)-2H-1-benzopyran-4-yl]phenylamino]acetic acid, ethyl ester;

(3R-trans)-3,4-dihydro-3-hydroxy-4-[N-[[3-(hydroxymethyl)-1,2,4-oxadiazol-5-yl)methyl]phenylamino]-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-[N-[3,4-dihydro-3-hydroxy-2,2-dimethyl-8-(trifluoromethyl)-2H-1-benzopyran-4-yl]phenylamino]acetic acid, ethyl ester;

(3R-trans)-4-[(4-chlorophenyl)[[3-(hydroxymethyl)-1,2,4-oxadiazol-5-yl)methyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(4-fluorophenyl)[[3-(hydroxymethyl)-1,2,4-oxadiazol-5-yl)methyl]amino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[N-[(3-amino-1,2,4-oxadiazol-5-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-[N-(6-benzoyl-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)phenylamino]acetic acid, ethyl ester;

(3R-trans)-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)[4-fluoro-3-[(phenylmethoxy)carbonyl]phenyl]amino]acetic acid, ethyl ester;

(3R-trans)-[[5-[(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)(4-fluorophenyl)amino]methyl]-1,2,4-oxadiazol-3-yl]methoxy]acetic acid;

(3R-trans)-[[5-[(4-chlorophenyl)(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)amino]methyl]-1,2,4-oxadiazol-3-yl]methoxy]acetic acid;

(3R-trans)-[(3-carboxy-4-fluorophenyl)(6-cyano-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-4-yl)amino]acetic acid, ethyl ester;

(3R-trans)-4-[(4-fluorophenyl)((2-methyl-2H-tetrazol-5-yl)-methylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[(4-fluorophenyl)((1-methyl-1H-tetrazol-5-yl)-methylamino)-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-(2-methylsulfinyl)ethyl]phenylamino]-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-(2-(methylsulfonyl)ethyl)phenylamino]-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[4-chloro-N-[(1H-imidazol-2-yl)methyl]-phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, monohydrochloride;

(3R-trans)-4-[4-chloro-N-[(1H-imidazol-2-yl)methyl]-phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, hydrogensulfate;

(3R-trans)-4-[4-chloro-N-[(1H-imidazol-2-yl)methyl]-phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, methanesulfonate;

(3R-trans)-4-[4-chloro-N-[(1H-imidazol-2-yl)methyl]-phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, phosphate;

(3R-trans)-4-[4-chloro-N-[(1H-imidazol-2-yl)methyl]-phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile, nitrate;

[3R-[3a,4b(R*)]]-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(5-oxo-1-(phenylmethyl)-2-pyrrolidinyl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile;

[3R-[3a,4b(S*)]]-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(5-oxo-2-pyrrolidinyl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile;

[3R-[3a,4a(R*)]]-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(5-oxo-2-pyrrolidinyl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-4-[N-[(1,5-dimethyl-1H-pyrazol-3-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

(3R-trans)-3,4-dihydro-3-hydroxy-2,2-dimethyl-4-[N-[(5-methyl-1H-pyrazol-3-yl)methyl]phenylamino]-2H-1-benzopyran-6-carbonitrile;


(3R-trans)-4-[N-[(1,3-dimethyl-1H-pyrazol-5-yl)methyl]phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile;

[3R-[3 α ,4B-(Z)]]-4-[N-(2-amino-4-oxo-2-pentenyl)phenylamino]-3,4-dihydro-3-hydroxy-2,2-dimethyl-2H-1-benzopyran-6-carbonitrile; or pharmaceutically acceptable salts thereof.

4. A pharmaceutical composition comprising a compound of Claim 1 and a pharmaceutically acceptable carrier.

5. A method for treating ischemia comprising administering to a mammalian specie in need thereof a therapeutically effective amount of a composition of Claim 4.

6. A compound of the formula




or pharmaceutically acceptable salts thereof wherein

a, b and d are all carbon atoms or one of a, b and d is a nitrogen atom or -N(O)- and the others are carbon atoms;


Y is a single bond, -CH₂-, -C(O)-, -O-, -S- or -N(R)- where R is hydrogen, alkyl, haloalkyl, aryl, arylalkyl, cycloalkyl or (cycloalkyl)alkyl;

R¹ is aryl or heterocyclo;

R^2 is $-\text{COOR}$, $-\text{CO-amino}$, $-\text{CO-substituted amino}$, amino, substituted amino, $-\text{NRCO-amino}$, $-\text{NRCO-substituted amino}$, $-\text{NRCOR}$, $-\text{NRSO}_2\text{R}$, $-\text{NR}(\text{C}=\text{NCN})\text{-amino}$, $-\text{NR}(\text{C}=\text{NCN})\text{-substituted amino}$.



$-\text{SR}$, $-\text{SOR}$, $-\text{SO}_2\text{R}$, $-\text{OR}$, cyano, heterocyclo, $-\text{CH}(\text{OR})_2$.




(where Z is O or H₂);

R^3 is hydrogen, hydroxy or $-\text{OC}(\text{O})\text{R}$;

R^4 and R^5 are each independently hydrogen, alkyl or arylalkyl, or R^4 and R^5 taken together with the carbon atom to which they are attached form a 5- to 7-membered carbocyclic ring;

R^6 is hydrogen, alkyl, haloalkyl, alkenyl, alkynyl, cycloalkyl, arylalkyl, (cycloalkyl)alkyl, $-\text{CN}$, $-\text{NO}_2$, $-\text{COR}$, $-\text{COOR}$,

$-\text{CONHR}$, $-\text{CON}(\text{R})_2$, $-\text{CF}_3$, $-\text{S-alkyl}$, $-\text{SOalkyl}$, $-\text{SO}_2\text{alkyl}$, $-\text{P}(\text{O-alkyl})_2$,



halogen, amino, substituted amino, $-\text{O-alkyl}$, $-\text{OCF}_3$, $-\text{OCH}_2\text{CF}_3$, $-\text{OCOalkyl}$, $-\text{OCONRalkyl}$, $-\text{NRCOalkyl}$, $-\text{NRCOOalkyl}$ or $-\text{NRCONR}$, tetrazolyl, imidazole, oxazole or triazole;

R^7 is hydrogen, alkyl, hydroxy, $-\text{O-alkyl}$, amino, substituted amino, $-\text{NHCOR}$, $-\text{CN}$ or $-\text{NO}_2$;

X is alkyl; or X- R^2 together are hydrogen, aryl or heterocyclo when R^1 is heterocyclo; and

n is an integer of 1 to 3.

7. A pharmaceutical composition comprising an effective amount of a compound of Claim 1 or 2, or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable carrier therefor.

8. A pharmaceutical composition comprising an effective amount of a compound of Claim 3, or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable carrier therefor.

9. A pharmaceutical composition for use in the treatment of ischemia in a patient comprising a therapeutically effective amount of a compound of Claim 1 or 2, or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable carrier therefor.

10. A pharmaceutical composition for use in the treatment of ischemia in a patient comprising a therapeutically effective amount of a compound of Claim 3, or a pharmaceutically acceptable salt thereof, together with a pharmaceutically acceptable carrier therefor.